

19 October 2006

1. Exercise Sheet Linear Algebra I for MCS Winter Term 2006/2007

(E1.1) [Geometric interpretation of vectors.]

- (i) Consider a (non-regular) hexagon $ABCDEF$. The following vertices and sides are given as vectors (points are given as vectors starting from the origin):

$$A = \begin{pmatrix} 0 \\ -2 \end{pmatrix} \quad D = \begin{pmatrix} 1 \\ 4 \end{pmatrix} \quad \overrightarrow{AB} = \begin{pmatrix} -2 \\ 2 \end{pmatrix} \quad \overrightarrow{BC} = \begin{pmatrix} 1 \\ 3 \end{pmatrix} \quad \overrightarrow{ED} = \begin{pmatrix} -3 \\ 3 \end{pmatrix} \quad \overrightarrow{FA} = \begin{pmatrix} -3 \\ -1 \end{pmatrix}$$

Here we write vectors as columns instead of rows, $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$ rather than (x_1, x_2) .

Can you draw the hexagon? Calculate \overrightarrow{AE} and compare with your drawing.

- (ii) Let $ABCD$ be an arbitrary quadrangle, where the vertices are given as vectors from the origin. Can you write down a condition that is true if and only if the quadrangle is a parallelogram?


(E1.2) [Parametric representation of a line.]

Consider the linear equation $E : ax_1 + bx_2 = c$ over \mathbb{R}^2 .

- (i) Let $a = 2$, $b = 3$ and $c = 6$. Find in that case the parametric representation for $S(E)$.
- (ii) Now in the general case, let $a \neq 0$. Find the parametric representation of $S(E)$.
- (iii) Consider lines in \mathbb{R}^3 .
Find a parametric representation of the line t through the points $(3, 0, 5)$ and $(1, 1, 1)$.

(E1.3) [Parametric representation.]

Draw a triangle ABC and consider the angle BAC with apex [*Scheitelpunkt*] A (we assume that the angle is strictly between 0 and π). Let P be the midpoint of AB and Q the midpoint of AC



- (i) Describe the lines through CP and through BQ using a parametric representation involving only the vectors $\mathbf{b} = \overrightarrow{AC}$ and $\mathbf{c} = \overrightarrow{AB}$ and the point A .
- (ii) Determine the intersection of the two lines from part(a).
- (iii) If you take B or C as apex instead of A , you can argue in a similar way. Which well-known theorem about triangles can you now prove?

(E1.4) [Intersections of planes.]

Consider three arbitrary planes P_1, P_2 and P_3 in \mathbb{R}^3 .

- (i) What are the possible intersections of P_1 and P_2 ?
- (ii) Now consider the intersection of P_1, P_2 and P_3 geometrically. Identify all possibilities and draw a picture for each case.
- (iii) In the case of an empty intersection of all three planes: which intersection types between pairs of planes can we have.