

December 14, 2006

## 9th Exercise Sheet Linear Algebra I for MCS Winter Term 2006/2007

### (E9.1) [Endomorphisms]

- (i) Let  $V$  be a vector space and  $\varphi : V \rightarrow V$  a vector space endomorphism. Prove that  $\text{image}(\varphi) \subseteq \ker(\varphi)$  if and only if  $\varphi \circ \varphi = \mathbf{0}$  (the null map, constant with value  $\mathbf{0}$ ).
- (ii) Find a linear map  $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that  $\ker(\varphi) = \text{image}(\varphi)$ .

### (E9.2) [Matrices for endomorphisms]

- (i) Consider the endomorphism of  $\mathbb{R}^2$  corresponding to the reflection in the line through  $\mathbf{0}$  that is spanned by the vector  $\mathbf{x} = (2, 1)$ . Obtain its matrix representation with respect to the standard basis.  
*Hint:* Draw a sketch and construct the images of the vectors  $(5, 0)$  and  $(0, 5)$ .
- (ii) Can you think of a basis of  $\mathbb{R}^2$  with respect to which the matrix looks much simpler? Can you generalize this to arbitrary reflections?
- (iii) Now consider the rotation in  $\mathbb{R}^3$  through 120 degrees about the axis through  $\mathbf{0}$  spanned by the vector  $\mathbf{x} = (1, 1, 1)$ . Draw a sketch of the situation and obtain the matrix representation with respect to a suitable basis.

### (E9.3) [Matrices and linear maps]

Let  $\varphi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a linear map such that  $\varphi(x, y, z) = (x + y + z, x + 2z, x + 2y)$ .

- (i) Determine  $\dim(\ker(\varphi))$ .

- (ii) Find bases  $B_1, B_2$  of  $\mathbb{R}^3$  such that 
$$[[\varphi]]_{B_2}^{B_1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

- (iii) Let  $\varphi : V \rightarrow V$  be an endomorphism of an arbitrary  $n$ -dimensional vector space. Prove that there exists bases  $B_1, B_2$  of  $V$ , such that

$$\llbracket \varphi \rrbracket_{B_2}^{B_1} = \left( \begin{array}{c|c} E_r & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} \end{array} \right),$$

where  $r = \dim(\text{image}(\varphi))$  and  $E_r$  is the  $r \times r$ -unit matrix.

**(E9.4) [An alternative proof of the dimension formula]**

- (i) Let  $\varphi : V \rightarrow W$  be a linear map. Prove that  $V/\ker(\varphi)$  is isomorphic to  $\text{image}(\varphi)$ .  
(ii) Use this result to give an alternative proof of the dimension formula.

**(E9.5) [The vector space  $\text{Hom}(V, W)$ ]**

(This is Exercise 3.2.1 in the notes.) Let  $\mathbb{F}$  be a field,  $V$  and  $W$   $\mathbb{F}$ -vector spaces with labelled bases  $(\mathbf{a}_1, \dots, \mathbf{a}_n)$ ,  $(\mathbf{b}_1, \dots, \mathbf{b}_m)$ , respectively. Consider the maps  $\varphi_{ij}$  for  $1 \leq i \leq m$ ,  $1 \leq j \leq n$  defined by

$$\varphi_{ij}: \begin{array}{ccc} V & \longrightarrow & W \\ \sum_{\ell=1}^n \lambda_{\ell} \mathbf{a}_{\ell} & \longmapsto & \lambda_j \mathbf{b}_i. \end{array}$$

- (i) Prove that the maps  $\varphi_{ij}$  are linear and pairwise distinct.  
(ii) Prove that the maps  $\varphi_{ij}$  are linearly independent and span  $\text{Hom}(V, W)$ . Use this to determine the dimension of  $\text{Hom}(V, W)$ .

Hint: For linear independence observe that a linear map (here: a linear combination of the  $\varphi_{ij}$ ) is the null map iff it has value  $\mathbf{0}$  on all arguments (equivalently: on all basis vectors). Similarly for a representation of an arbitrary  $\varphi \in \text{Hom}(V, W)$  as a linear combination of the  $\varphi_{ij}$ , use that two linear maps are equal if they agree on all basis vectors. The rest is matching of coefficients.