

January 11, 2006

## 11th Exercise Sheet Linear Algebra I for MCS Winter Term 2006/2007

### (E11.1) [Zero divisors]

- (i) Let  $A \in \mathbb{F}^{(n,n)}$  be some matrix. Show that  $A$  is not invertible if, and only if, there exists some matrix  $B \in \mathbb{F}^{(n,n)}$ ,  $B \neq \mathbf{0}$  with  $AB = \mathbf{0}$ .  
 (*Hint*: use the fact that a matrix represents a linear map and choose a suitable basis.)
- (ii) (\*) Let now  $R$  be a ring and  $a \in R$ . Which directions of the statement “ $a$  is not invertible if, and only if, there exists some  $b \in R$  with  $ab = 0$ .” still hold? Give proof(s) or counterexample(s).

### (E11.2) [Calculating with block matrices]

- (i) Let  $A_{11} \in \mathbb{F}^{(n,p)}$ ,  $A_{12} \in \mathbb{F}^{(n,q)}$ ,  $B_{11} \in \mathbb{F}^{(p,m)}$  and  $B_{21} \in \mathbb{F}^{(q,m)}$ . Show that the product of the block matrices can be calculated as

$$\begin{bmatrix} A_{11} & A_{12} \end{bmatrix} \cdot \begin{bmatrix} B_{11} \\ B_{21} \end{bmatrix} = \begin{bmatrix} A_{11}B_{11} + A_{12}B_{21} \end{bmatrix}.$$

What is the size of the resulting matrix?

- (ii) Let  $A$  and  $B$  be two  $n \times n$  block matrices of the form

$$A = \begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix}, \quad B = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix},$$

where  $A_{11}, B_{11} \in \mathbb{F}^{(p,p)}$ ,  $A_{22}, B_{22} \in \mathbb{F}^{(n-p,n-p)}$ , etc.

Calculate the product of  $A$  and  $B$  in blocks:

$$\begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix} \cdot \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} ? & ? \\ ? & ? \end{bmatrix}.$$

**(E11.3) [Projections with respect to a suitable basis]**

Let  $V = U \oplus W$  be a finite dimensional vector space, and  $\pi : V \rightarrow V$  a projection (i.e.  $\pi \circ \pi = \pi$ , cf. T9.2) such that  $U = \ker(\pi)$  and  $W = \text{image}(\pi)$ .

- (i) Show that there exists a basis of  $V$  such that the matrix of  $\pi$  with respect to that basis is a diagonal matrix with only ones and zeroes on the diagonal.<sup>1</sup>

*Hint:* start from bases of  $U$  and  $W$ , respectively.

- (ii) Let  $\pi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be given by the matrix

$$A = \begin{pmatrix} 0 & -1 & 1 \\ 4 & 5 & -4 \\ 4 & 4 & -3 \end{pmatrix}$$

with respect to the standard basis. Verify that  $\pi$  is a projection. Determine a matrix  $S$  and a diagonal matrix  $D$  such that  $D = S^{-1}AS$ .

**(E11.4) [Basis transformations]**

- (i) The linear map  $\varphi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is given by the matrix

$$[[\varphi]_E^E = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 3 & -1 \\ 1 & 1 & 1 \end{pmatrix}$$

w.r.t. the standard basis  $E$ .

Compute the matrix  $[[\varphi]_B^B$  of  $\varphi$  w.r.t. the basis  $B = ((1, -1, -1), (1, 0, 1), (1, 1, 1))$ .

- (ii) We consider the  $\mathbb{R}$ -vector spaces  $\mathbb{R}^2$  und  $\mathbb{R}^3$  with bases  $B = ((0, 1), (1, 0))$  and  $C = ((0, 1, 1), (1, 0, 1), (1, 1, 0))$ . A linear map  $\psi \in \text{Hom}(\mathbb{R}^3, \mathbb{R}^2)$  is given by

$$[[\psi]_B^C := \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}.$$

Derive the matrix of  $\psi$  w.r.t. the standard bases  $S_2, S_3$ .

---

<sup>1</sup>Compare this **carefully** to the representation of an arbitrary endomorphism in E9.3(iii). What are the similarities/differences?