

February 8, 2007

14th Exercise Sheet Linear Algebra I for MCS Winter Term 2006/2007

(E14.1) [Scalar product]

The *standard scalar product* [Standardskalarprodukt] on \mathbb{F}^n is defined to be the map

$$\langle \cdot, \cdot \rangle : \mathbb{F}^n \times \mathbb{F}^n \rightarrow \mathbb{F}, \quad (\mathbf{x}, \mathbf{y}) \mapsto \langle \mathbf{x}, \mathbf{y} \rangle := \sum_{i=1}^n x_i y_i.$$

Show that $\langle \cdot, \cdot \rangle$ has the following properties:

- (i) it is bilinear,
- (ii) it is symmetric (i.e. $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$),
- (iii) $\langle \mathbf{x}, \mathbf{e}_i \rangle = x_i$ for $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{F}^n$ and $1 \leq i \leq n$.

Two vectors \mathbf{x} and \mathbf{y} are called *orthogonal* [orthogonal], $\mathbf{x} \perp \mathbf{y}$, if $\langle \mathbf{x}, \mathbf{y} \rangle = 0$.

(E14.2) [Cross product]

Let \mathbb{F}^3 be equipped with the standard scalar product (as above). Consider a function

$$\times : \mathbb{F}^3 \times \mathbb{F}^3 \rightarrow \mathbb{F}^3, \quad (\mathbf{x}, \mathbf{y}) \mapsto \mathbf{x} \times \mathbf{y},$$

with the following properties:

- bi-linearity (linearity in each argument, as an \mathbb{F}^3 -valued function),
- antisymmetry (i.e., for all $\mathbf{x} \in \mathbb{F}^3$, $\mathbf{x} \times \mathbf{x} = \mathbf{0}$),
- $\mathbf{x} \times \mathbf{y}$ is orthogonal to \mathbf{x} and \mathbf{y} ,
- normalization: $\mathbf{e}_1 \times \mathbf{e}_2 = \mathbf{e}_3$.

The aim of this exercise is to show that a function with these properties is uniquely determined, and to give an explicit description of this function.

- (i) Prove that the function $g : \mathbb{F}^3 \times \mathbb{F}^3 \times \mathbb{F}^3 \rightarrow \mathbb{F}$, $(\mathbf{x}, \mathbf{y}, \mathbf{z}) \mapsto \langle \mathbf{x} \times \mathbf{y}, \mathbf{z} \rangle$ is 3-linear, antisymmetric and normalized (in the sense that $g(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3) = 1$).
- (ii) Conclude that g and hence \times is uniquely determined.
- (iii) Use E14.1(iii) to give an explicit formula for $\mathbf{x} \times \mathbf{y}$ in terms of x_i, y_i , and show that the function thus obtained satisfies all requirements.
- (iv) Show that \times is not associative.

(E14.3) [Symmetries of patterns]

Consider a subgroup H of the group of affine transformations of \mathbb{R}^2 (cf. section 3.4 of the notes) and let

$$H_0 := \{\varphi \in \text{GL}_2(\mathbb{R}) : [\varphi, \mathbf{u}] \in H \text{ for some } \mathbf{u} \in \mathbb{R}^2\} \subseteq \text{GL}_2(\mathbb{R});$$

$$L_H := \{\mathbf{u} \in \mathbb{R}^2 : \tau_{\mathbf{u}} \in H\} \subseteq \mathbb{R}^2.$$

- (i) Show that H_0 is a subgroup of $\text{GL}_2(\mathbb{R})$. Give an example that not necessarily $H_0 \subseteq H$.
- (ii) Show that L_H is closed under vector addition (a subgroup of $(\mathbb{R}^2, +, \mathbf{0})$) and that L_H is mapped onto itself by H_0 .
- (iii) (*) Assuming that $L_H \neq \{\mathbf{0}\}$ does not contain arbitrarily small non-zero vectors, show that L_H is the set of integer linear combinations of one or two vectors in $\mathbb{R}^2 \setminus \{\mathbf{0}\}$ (a one- or two-dimensional lattice).
- (iv) Try to describe the symmetry group H of the pattern below as a subgroup of the group of affine transformations. Determine L_H and H_0 and try to describe how these fit together in H . Can you find a small finite set of transformations in H that generate H in the sense that all elements of H are compositions of the given ones?

