

November 7, 2006

4th Tutorial Sheet Linear Algebra I for MCS Winter Term 2006/2007

(T4.1) Function spaces

Let A be any set, \mathbb{F} a field. Let $\mathcal{F}(A, \mathbb{F})$ be the set of all functions $f : A \rightarrow \mathbb{F}$. Show that $\mathcal{F}(A, \mathbb{F})$ with point-wise addition and point-wise multiplication is an \mathbb{F} -vector space.

(A borderline case: is it ok to admit $A = \emptyset$?)

(T4.2) Vector spaces of polynomial functions

Let V be the set $\text{Pol}_7(\mathbb{R})$, the space of polynomial functions of degree 7 over \mathbb{R} .

Show that V is an \mathbb{R} -vector space with point-wise addition and point-wise multiplication.

Determine whether or not W is a subspace of V where

- (i) W consists of the degree 7 polynomials with integral coefficients.
- (ii) W consists of the degree 7 polynomials with only even powers of x of the form $b_0 + b_1x^2 + b_2x^4 + b_3x^6$.

(T4.3) Polynomials vs polynomial functions

Consider a finite field \mathbb{F}_p . Two distinct polynomials over \mathbb{F}_p , $q(x) = \sum_{i=0}^n a_i x^i$ and $r(x) = \sum_{i=0}^n b_i x^i$ with $(a_0, \dots, a_n) \neq (b_0, \dots, b_n)$ may induce the same polynomial function.

- (i) Give concrete examples of this phenomenon, e.g., over \mathbb{F}_2 .
- (ii) Give a combinatorial counting argument why this must occur for sufficiently large degrees n .
- (iii) How is this different for the field \mathbb{R} ?