

Some constructions of modular forms for the Weil representation of $SL_2(\mathbb{Z})$

Nils R. Scheithauer,
Fachbereich Mathematik, Technische Universität Darmstadt,
Schloßgartenstraße 7, 64289 Darmstadt, Deutschland
scheithauer@mathematik.tu-darmstadt.de

Modular forms for the Weil representation of $SL_2(\mathbb{Z})$ play an important role in the theory of automorphic forms on orthogonal groups. In this paper we give some explicit constructions of these functions. As an application we construct new examples of generalized Kac-Moody algebras whose denominator identities are holomorphic automorphic products of singular weight. They correspond naturally to the Niemeier lattices with root systems D_{12}^2 , E_8^3 and to the Leech lattice.

- 1 Introduction
- 2 Modular forms for the Weil representation of $SL_2(\mathbb{Z})$
- 3 Liftings from congruence subgroups
- 4 Induction from isotropic subgroups
- 5 Discriminant forms of squarefree level
- 6 Some new generalized Kac-Moody algebras

1 Introduction

The singular theta correspondence (cf. [B2] and also [Br]) is a map from modular forms for the Weil representation of $SL_2(\mathbb{Z})$ to automorphic forms on orthogonal groups. More precisely let L be an even lattice of signature $(n, 2)$, $n > 2$ even with discriminant form D and F a modular form for the Weil representation of $SL_2(\mathbb{Z})$ on $\mathbb{C}[D]$ of weight $(2-n)/2$ which is holomorphic on the upper halfplane and has integral principal part. Then the integral

$$\int_{\mathcal{F}} F(\tau) \bar{\theta}(Z, \tau) y \frac{dx dy}{y^2},$$

where \mathcal{F} is the standard fundamental domain of $SL_2(\mathbb{Z})$ on the upper halfplane and θ is the Siegel theta function of L , can be regularized. Let $\Psi(Z)$ be the exponential of the regularized value. Then $\Psi(Z)$ is an automorphic form for a subgroup of $O(L)$ which has nice product expansions at the rational 0-dimensional cusps. The function $\Psi(Z)$ is called the automorphic product associated to F . The weight of Ψ is determined by the constant coefficient of F_0 and its divisor by the principal part of F .

The smallest possible weight of a nonconstant holomorphic automorphic form on $O_{n,2}(\mathbb{R})$ is given by $(n-2)/2$. Holomorphic automorphic forms of this so-called singular weight are particularly interesting because their Fourier coefficients are supported only on isotropic vectors. It has turned out that the denominator identities of infinite-dimensional Lie algebras are sometimes holomorphic automorphic products of singular weight. These Lie algebras describe strings moving on suitable orbifolds and they seem to be very rare [S1]. So far there are only 12 known examples [S2].

In this paper we give some explicit constructions of modular forms for the Weil representation of $SL_2(\mathbb{Z})$. As an application we construct 3 new generalized Kac-Moody algebras whose denominator identities are holomorphic automorphic products of singular weight. In contrast to the previous examples the corresponding vector valued modular forms are not symmetric under $O(D)$ which makes their construction more difficult.

We describe the results in more detail.

First we show how scalar valued modular forms on congruence subgroups induce modular forms for the Weil representation of $SL_2(\mathbb{Z})$. For $\Gamma_1(N)$ the result is as follows (see Theorems 3.1 and 3.7).

Let D be a discriminant form of even signature and level dividing N and ρ_D the Weil representation of $SL_2(\mathbb{Z})$ on $\mathbb{C}[D]$. Let $\gamma \in D$ and f be a modular form on $\Gamma_1(N)$ with character χ_γ . Then

$$F_{\Gamma_1(N),f,\gamma} = \sum_{M \in \Gamma_1(N) \backslash \Gamma} f|_M \rho_D(M^{-1}) e^\gamma$$

is a modular form for ρ_D . Every modular form for ρ_D is a linear combination of such functions. The function $F_{\Gamma_1(N),f,\gamma}$ can be written as a sum $\sum F_s$ over the cusps of $\Gamma_1(N)$ where

$$F_s = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\mu \in a\gamma + D^{c*}} e(d(\mu - a\gamma)_c^2/2) e(b\mu\gamma) e(-ab\gamma^2/2) \\ t g_{mt,j\mu} \{ e^\mu + (-1)^k e(\text{sign}(D)/4) e^{-\mu} \}.$$

if $N > 2$ and s is regular and similarly in the other cases.

There are analogous results for the congruence subgroups $\Gamma(N)$ and $\Gamma_0(N)$. However the liftings are not equivalent. In section 6 we give an example of a modular form for the Weil representation which is not induced from $\Gamma_0(N)$.

Next we describe an induction from the isotropic subgroups of a discriminant form (cf. Theorem 4.1).

Let D be a discriminant form of even signature and H an isotropic subgroup of D . Let $F_{D_H} = \sum_{\gamma \in D_H} F_{D_H, \gamma} e^\gamma$ be a modular form for the Weil representation of the discriminant form $D_H = H^\perp/H$. Then

$$F = \sum_{\gamma \in H^\perp} F_{D_H, \gamma+H} e^\gamma$$

is a modular form for ρ_D .

Modular forms which are induced in this way can be considered as oldforms.

Let D be a discriminant form of even signature and level dividing N . We have seen that all modular forms for ρ_D are induced from $\Gamma(N)$ or $\Gamma_1(N)$. However calculating the corresponding liftings is often laborious because the indices of these groups in $SL_2(\mathbb{Z})$ are rather large. We give a natural sufficient condition for a modular form for ρ_D to be induced from $\Gamma_0(N)$ (see Proposition 5.3 and Theorem 5.4).

Let D be a discriminant form of squarefree level N and $F = \sum F_\gamma e^\gamma$ a modular form for ρ_D which is invariant under $O(D)$. Then the complex vector space W spanned by the components F_γ , $\gamma \in D$ is generated by the functions $F_0|_M$, $M \in SL_2(\mathbb{Z})$. Let W_0 be the subspace of W with T -eigenvalue $e(0)$. Then the map

$$\begin{aligned} \Phi: W_0 &\longrightarrow W_0 \\ f &\longmapsto 0\text{-component of } F_{\Gamma_0(N), f, 0} \end{aligned}$$

is a bijection. In particular $F = F_{\Gamma_0(N), f, 0}$ for a suitable function f in W_0 .

Finally we use the above results to construct some modular forms for the Weil representation with nonnegative integral coefficients and reflective poles. The theta lifts of these functions have singular weight and give the denominator identities of some new generalized Kac-Moody algebras. The simplest example is the following (see section 6).

Let N be the Niemeier lattice with root system E_8^3 . Let g be a permutation of the three E_8 -components of order 3. Then the fixed point sublattice N^g of g is isomorphic to $\sqrt{3}E_8$ and the orthogonal complement N^{g^\perp} is isomorphic to $A_2 \otimes E_8$. The theta function $\theta_{N^{g^\perp}}$ defines a modular form for the discriminant form of N^g . This function is invariant under $O(N^g)$ because the centralizer of g in $O(N)$ induces the full orthogonal group of N^g . Let $L = N^g \oplus \begin{pmatrix} -2 & 3 \\ 3 & 0 \end{pmatrix}$. Then $\theta_{N^{g^\perp}}$ induces a modular form on L . Denote the quotient of this form by the invariant 3Δ by $F_{\theta_{N^{g^\perp}/3\Delta}}$. Define $\eta_g(\tau) = \eta(3\tau)^8$. Then the sum

$$F = F_{\theta_{N^{g^\perp}/3\Delta}} + \frac{1}{3} F_{\Gamma_0(9), 1/\eta_g, 0}$$

is a modular form for the Weil representation of $L \oplus II_{1,1}$ with nonnegative integral coefficients and reflective poles. The theta lift of F is holomorphic, has singular weight and is up to a constant given by

$$\begin{aligned} e((\rho, Z)) \prod_{\alpha \in L'^+} (1 - e((\alpha, Z)))^{[F_{\alpha+L}](-\alpha^2/2)} \\ = \sum_{w \in W} \det(w) e((w\rho, Z)) \prod_{n>0} (1 - e((3nw\rho, Z)))^8. \end{aligned}$$

This is the denominator identity of a generalized Kac-Moody algebra with root lattice L' whose multiplicities and simple roots can be described easily.

There are similar examples for the Niemeier lattice with root system D_{12}^2 and for the Leech lattice.

The paper is organized as follows.

In section 2 we recall some properties of modular forms for the Weil representation of $SL_2(\mathbb{Z})$.

Then we describe the liftings from modular forms on congruence subgroups to modular forms for the Weil representation of $SL_2(\mathbb{Z})$.

In the next section we show that a modular form for the Weil representation of an even lattice induces a modular form for the Weil representation of any finite index sublattice.

In section 5 we show that symmetric modular forms on discriminant forms of squarefree level N are induced from $\Gamma_0(N)$.

Finally we use the above methods to construct some new generalized Kac-Moody algebras with automorphic denominator identity. These Lie algebras correspond naturally to the Niemeier lattices with root systems D_{12}^2 and E_8^3 and to the Leech lattice.

The author thanks R. E. Borcherds, G. Höhn and F. Werner for stimulating discussions and helpful comments.

2 Modular forms for the Weil representation of $SL_2(\mathbb{Z})$

In this section we recall some properties of modular forms for the Weil representation of $SL_2(\mathbb{Z})$ from [S2].

Let D be a discriminant form with quadratic form $D \rightarrow \mathbb{Q}/\mathbb{Z}, \gamma \mapsto \gamma^2/2$ (cf. also [N], [CS]). The level of D is the smallest positive integer N such that $N\gamma^2/2 = 0 \pmod 1$ for all $\gamma \in D$.

Let c be an integer. Then c acts by multiplication on D and we have an exact sequence $0 \rightarrow D_c \rightarrow D \rightarrow D^c \rightarrow 0$ where D_c is the kernel and D^c the image of this map. The group D^c is the orthogonal complement of D_c .

Let D^{c*} be the set of elements $\alpha \in D$ satisfying $c\gamma^2/2 + \alpha\gamma = 0 \pmod 1$ for all $\gamma \in D_c$. Then D^{c*} is a coset of D^c . When we have chosen a Jordan decomposition of D then there is a canonical coset representative x_c of D^{c*} satisfying $2x_c = 0$. Write $\alpha \in D^{c*}$ as $\alpha = x_c + c\gamma$. Then $\alpha_c^2/2 = c\gamma^2/2 + x_c\gamma \pmod 1$ is independent of the choice of γ . This gives a well defined map $D^{c*} \rightarrow \mathbb{Q}/\mathbb{Z}, \alpha \mapsto \alpha_c^2/2$.

More generally let H be an isotropic subgroup of D . We define D_H^{c*} as the set of elements $\alpha \in D$ satisfying $c\gamma^2/2 + \alpha\gamma = 0 \pmod 1$ for all $\gamma \in c^{-1}(H) \cap H^\perp$. Then $D_H^{c*} = D^{c*}$ if $H = 0$.

Suppose D has even signature. We define a scalar product on the group ring $\mathbb{C}[D]$ which is linear in the first and antilinear in the second variable by $(e^\gamma, e^\beta) = \delta^{\gamma\beta}$. Then there is a unitary action of the group $\Gamma = SL_2(\mathbb{Z})$ on $\mathbb{C}[D]$

defined by

$$\begin{aligned}\rho_D(T)e^\gamma &= e(-\gamma^2/2) e^\gamma \\ \rho_D(S)e^\gamma &= \frac{e(\text{sign}(D)/8)}{\sqrt{|D|}} \sum_{\beta \in D} e(\gamma\beta) e^\beta\end{aligned}$$

where $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $T = \begin{pmatrix} 1 & \\ 0 & 1 \end{pmatrix}$ are the standard generators of Γ . This representation is called the Weil representation of Γ on $\mathbb{C}[D]$. It commutes with $O(D)$.

Let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Then

$$\rho_D(M)e^\gamma = \xi \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in D^{c*}} e(-a\beta_c^2/2) e(-b\beta\gamma) e(-bd\gamma^2/2) e^{d\gamma+\beta}$$

where $\xi = e(\text{sign}(D)/4) \prod \xi_p$. The local factors ξ_p can be expressed by means of the Jordan components of D (cf. [S2]).

Let

$$F(\tau) = \sum_{\gamma \in D} F_\gamma(\tau) e^\gamma$$

be a holomorphic function on the upper halfplane with values in $\mathbb{C}[D]$ and k an integer. Then F is a modular form for ρ_D of weight k if

$$F(M\tau) = (c\tau + d)^k \rho_D(M) F(\tau)$$

for all $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in Γ and F is meromorphic at ∞ . Note that the transformation formula can be written equivalently as

$$F = \sum_{\gamma \in D} F_\gamma|_{M^{-1}} \rho_D(M) e^\gamma$$

for all $M \in \Gamma$.

The components of a modular form $F = \sum_{\gamma \in D} F_\gamma e^\gamma$ for ρ_D transform as

$$F_\gamma|_M = \bar{\xi}(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in D^{c*}} e(-d\beta_c^2/2) e(-b\beta\gamma) e(-ab\gamma^2/2) F_{a\gamma+\beta}$$

where $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ and $\xi(M^{-1})$ is the above root of unity corresponding to M^{-1} .

Classical examples of modular forms transforming under the Weil representation are theta functions. Let L be a positive definite even lattice of even rank $2k$ with discriminant form D . For $\gamma \in D$ define

$$\theta_\gamma(\tau) = \sum_{\alpha \in \gamma + L} q^{\alpha^2/2}$$

with $q^{\alpha^2/2} = e(\tau\alpha^2/2)$. Then

$$\theta = \sum_{\gamma \in D} \theta_\gamma e^\gamma$$

is a modular form for the dual Weil representation $\bar{\rho}_D$ of weight k which is holomorphic at ∞ .

Let D and D' be two discriminant forms of even signature and let $i : D \rightarrow D'$ be an isomorphism of groups satisfying $i(\gamma)^2/2 = -\gamma^2/2$ for all $\gamma \in D$. If $F_{D'} = \sum_{\gamma \in D'} F_\gamma e^\gamma$ is a modular form for the dual Weil representation $\bar{\rho}_{D'}$, then $F_D = \sum_{\gamma \in D} F_{i(\gamma)} e^\gamma$ transforms under ρ_D . For example if L is an even unimodular lattice and K a primitive sublattice of L with orthogonal complement K^\perp then we have a natural isomorphism $i : D_K \rightarrow D_{K^\perp}$ such that $i(\gamma)^2/2 = -\gamma^2/2$.

3 Liftings from congruence subgroups

In this section we describe how modular forms on congruence subgroups induce modular forms on Γ transforming under the Weil representation and we calculate these liftings explicitly. For the congruence subgroup $\Gamma_0(N)$ these results are already known [S2]. We include them here because we need the corresponding formulas later.

Let D be a discriminant form of even signature and N a positive integer such that the level of D divides N . Let $F = \sum_{\gamma \in D} F_\gamma e^\gamma$ be a modular form for ρ_D and $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$. Then the formula for ρ_D gives

$$\rho_D(M)e^\gamma = \left(\frac{a}{|D|} \right) e((a-1)\text{oddy}(D)/8) e(-bd\gamma^2/2) e^{d\gamma}$$

and

$$F_\gamma|_M = \left(\frac{d}{|D|} \right) e((d-1)\text{oddy}(D)/8) e(-ab\gamma^2/2) F_{a\gamma}.$$

We define the quadratic Dirichlet character $\chi_D : \Gamma_0(N) \rightarrow \mathbb{C}^*$ by

$$\chi_D \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) = \left(\frac{a}{|D|} \right) e((a-1)\text{oddy}(D)/8)$$

and for $\gamma \in D$ the character $\chi_\gamma : \Gamma_1(N) \rightarrow \mathbb{C}^*$ with

$$\chi_\gamma \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) = e(-b\gamma^2/2).$$

Then F_0 is a modular form on $\Gamma_0(N)$ of character χ_D and F_γ is a modular form on $\Gamma_1(N)$ with character χ_γ . Conversely we have the following result.

Theorem 3.1

Let D be a discriminant form of even signature and level dividing N .

1. Let f be a scalar valued modular form on $\Gamma_0(N)$ of weight k and character χ_D and H an isotropic subset of D which is invariant under $(\mathbb{Z}/N\mathbb{Z})^*$ as a set. Then

$$F_{\Gamma_0(N),f,H} = \sum_{M \in \Gamma_0(N) \setminus \Gamma} \sum_{\gamma \in H} f|_M \rho_D(M^{-1}) e^\gamma$$

is a modular form for ρ_D of weight k which is invariant under the automorphisms of the discriminant form that stabilize H as a set.

2. Let $\gamma \in D$ and f a scalar valued modular form for $\Gamma_1(N)$ of weight k and character χ_γ . Then

$$F_{\Gamma_1(N),f,\gamma} = \sum_{M \in \Gamma_1(N) \setminus \Gamma} f|_M \rho_D(M^{-1}) e^\gamma$$

is a modular form for ρ_D of weight k which is invariant under the stabilizer of γ in $O(D)$.

3. Finally if f is a scalar valued modular form on $\Gamma(N)$ of weight k and $\gamma \in D$ then

$$F_{\Gamma(N),f,\gamma} = \sum_{M \in \Gamma(N) \setminus \Gamma} f|_M \rho_D(M^{-1}) e^\gamma$$

is a modular form for ρ_D of weight k which is invariant under the stabilizer of γ in $O(D)$.

Proof: We have to show that the liftings are well defined, transform correctly under Γ and have the stated symmetries. We describe this in the second case. The other cases are analogous.

For $M \in \Gamma$ we define the function $F_M = f|_M \rho_D(M^{-1}) e^\gamma$. Then

$$F_{KM} = f|_{KM} \rho_D((KM)^{-1}) e^\gamma = \chi_\gamma(K) f|_M \rho_D(M^{-1}) \rho_D(K^{-1}) e^\gamma = F_M$$

for $K \in \Gamma_1(N)$. Hence $F = \sum_{M \in \Gamma_1(N) \setminus \Gamma} F_M$ is well defined. Now let $K = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Then

$$\begin{aligned} F(K\tau) &= \sum_{M \in \Gamma_1(N) \setminus \Gamma} f|_M(K\tau) \rho_D(M^{-1}) e^\gamma \\ &= (c\tau + d)^k \sum_{M \in \Gamma_1(N) \setminus \Gamma} f|_{MK}(\tau) \rho_D(M^{-1}) e^\gamma \\ &= (c\tau + d)^k \rho_D(K) \sum_{M \in \Gamma_1(N) \setminus \Gamma} f|_{MK}(\tau) \rho_D(K^{-1}) \rho_D(M^{-1}) e^\gamma \\ &= (c\tau + d)^k \rho_D(K) \sum_{M \in \Gamma_1(N) \setminus \Gamma} f|_{MK}(\tau) \rho_D((MK)^{-1}) e^\gamma \\ &= (c\tau + d)^k \rho_D(K) F(\tau) \end{aligned}$$

by shifting the summation index. Finally the functions F_M and F are invariant under the stabilizer of γ in $O(D)$ because the Weil representation commutes with $O(D)$. \square

The liftings satisfy some obvious relations. However they are not equivalent. We will see that there are modular forms for the Weil representation which are not induced from $\Gamma_0(N)$. On the other hand let $F = \sum_{\gamma \in D} F_\gamma e^\gamma$ be a modular form for ρ_D . Then

$$\begin{aligned} F &= \frac{1}{|\Gamma/\Gamma_1(N)|} \sum_{M \in \Gamma_1(N) \backslash \Gamma} \sum_{\gamma \in D} F_\gamma |_{M} \rho_D(M^{-1}) e^\gamma \\ &= \frac{1}{|\Gamma/\Gamma_1(N)|} \sum_{\gamma \in D} F_{\Gamma_1(N), F_\gamma, \gamma}. \end{aligned}$$

Hence F can be written as a linear combination of liftings from $\Gamma_1(N)$. Finally let V be the complex vector space generated by the components F_γ . Then Γ acts on $V \otimes \mathbb{C}[D]$ by $\rho(M)(f \otimes e^\gamma) = f|_{M^{-1}} \otimes \rho_D(M) e^\gamma$. The invariants $(V \otimes \mathbb{C}[D])^\Gamma$ are modular forms for ρ_D . This space is spanned by the functions $F_{\Gamma(N), F_\gamma, \mu}$.

We calculate the liftings explicitly.

The group $\Gamma_0(N)$ has index $N \prod_{p|N} (1 + 1/p)$ in Γ and $\sum_{c|N} \phi(c, N/c)$ classes of cusps. Let $a/c \in \mathbb{Q}$ with $(a, c) = 1$. Then the equivalence class of a/c as a cusp of $\Gamma_0(N)$ is determined by the invariants (c, N) (a divisor of N) and $ac/(c, N)$ (a unit in $\mathbb{Z}/(c, N/(c, N))\mathbb{Z}$). The width of a/c is $t = N/(c^2, N)$ and the stabilizer of a/c in $\Gamma_0(N)$ is given by $\Gamma_0(N)_{a/c} = \{\pm T_{a/c}^n \mid n \in \mathbb{Z}\}$ with

$$T_{a/c} = MT^t M^{-1} = \begin{pmatrix} 1 - act & a^2 t \\ -c^2 t & 1 + act \end{pmatrix}$$

for any matrix $M \in \Gamma$ satisfying $M\infty = a/c$.

Let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Then the cosets of $\Gamma_0(N) \backslash \Gamma$ sending ∞ to a/c are given by MT^n where n ranges over a complete set of residues modulo $t = N/(c^2, N)$.

Let D be a discriminant form of even signature and level dividing N and f a modular form on $\Gamma_0(N)$ with character χ_D .

We denote the order of $\chi_D(T_{a/c})$ by m .

For $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ the function $f|_M$ has a Fourier expansion of the form

$$f|_M(\tau) = \sum a_n q_{mt}^n$$

with $q_{mt} = e(\tau/mt)$ and n integral if $m = 1$ and n odd integral if $m = 2$.

Let $a/c \in \mathbb{Q}$ with $(a, c) = 1$ and $\mu \in D^{c*}$. Then $\mu^2/2 = -n/mt \pmod{1}$ where n is integral if $m = 1$ and n odd integral if $m = 2$.

Let H be an isotropic subset of D which is invariant under $(\mathbb{Z}/N\mathbb{Z})^*$. Then we can write $\sum_{\gamma \in H} e^\gamma$ as a linear combination of sums $\sum_{\gamma \in I} e^\gamma$ where I is an isotropic subgroup of D . Therefore it is sufficient to calculate the lift F of f on H in the case that H is an isotropic subgroup. We will assume this in the following.

We can write F as

$$F = \sum_{s \in \Gamma_0(N) \backslash P} F_s$$

where P is the set of cusps of Γ and

$$F_s = \sum_{\substack{M \in \Gamma_0(N) \setminus \Gamma \\ M\infty = s}} \sum_{\gamma \in H} f|_M \rho_D(M^{-1}) e^\gamma.$$

The function F_s is T -invariant, i.e. $F_s(T\tau) = \rho_D(T)F_s(\tau)$. We have

Theorem 3.2

Let $a/c \in \mathbb{Q}$ be a representative of s with $(a, N) = 1$ and $c|N$ and $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Decompose $f|_M = g_{mt,0} + \dots + g_{mt,mt-1}$ into T -invariant functions $g_{mt,j}$ satisfying $g_{mt,j}|_T = e(j/mt)g_{mt,j}$ and for $w \in D^{c*}$ define j_w by $w^2/2 = -j_w/mt \pmod{1}$. Then

$$F_s = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} |H \cap cH^\perp| \sum_{v \in H/(H \cap D^c)} \sum_{w \in (D^{c*} \cap D_H^{c*})} e(dw_c^2/2) \Phi_{H,a,c}(w) t g_{mt,j_w} e^{v+w}$$

where

$$\Phi_{H,a,c}(w) = \sum_{\gamma \in G_{H,c}} e(ac\gamma^2/2 + w\gamma)$$

and

$$G_{H,c} = \frac{c^{-1}(H)/D_c}{((c^{-1}(H) \cap H^\perp) + D_c)/D_c}.$$

The group $\Gamma_1(N)$ has index $N^2 \prod_{p|N} (1 - 1/p^2)$ in Γ and

$$(1/2) \sum_{d|N} \phi(d)\phi(N/d) \begin{cases} 2 & \text{if } N = 2, \\ 3 & \text{if } N = 4, \\ \text{if } N = 3 \text{ or } N > 4 \end{cases}$$

classes of cusps. Two cusps a/c and a'/c' in \mathbb{Q} with $(a, c) = (a', c') = 1$ are equivalent modulo $\Gamma_1(N)$ if and only if

$$\begin{pmatrix} a' \\ c' \end{pmatrix} = \pm \begin{pmatrix} a + jc \\ c \end{pmatrix} \pmod{N}$$

for some j (cf. [DS], section 3.8). The cusp $1/2$ of $\Gamma_1(4)$ is irregular and has width $t = 1$. All other classes of cusps a/c are regular and have width $t = N/(c, N)$. For a cusp a/c in \mathbb{Q} with $(a, c) = 1$ and width t define as above $T_{a/c} = MT^t M^{-1}$ where M is any matrix in Γ sending ∞ to a/c . Then the stabilizer of a/c in $\Gamma_1(N)$ is given by $\{T_{a/c}^n \mid n \in \mathbb{Z}\}$ if a/c is regular and by $\{(-T_{a/c})^n \mid n \in \mathbb{Z}\}$ if a/c is irregular.

It is easy to prove the following result.

Proposition 3.3

Let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Then the cosets of $\Gamma_1(N) \backslash \Gamma$ sending ∞ to a/c are represented by

$$\begin{aligned} MT^n & \quad \text{if } N = 2, \\ \pm MT^n & \quad \text{if } N > 2 \end{aligned}$$

where n ranges over a complete set of residues modulo t .

Let D be a discriminant form of even signature and level dividing N and $\gamma \in D$ with $\gamma^2/2 = -j/N \pmod{1}$.

Proposition 3.4

Let $a/c \in \mathbb{Q}$ with $(a, c) = 1$ be a cusp of $\Gamma_1(N)$.

If a/c is regular then $\chi_\gamma(T_{a/c})$ has order $m = (c, N)/(c, N, j)$.

If a/c is irregular then $\chi_\gamma(-T_{a/c})$ has order $m = 4/(4, j)$.

Let f be a modular form on $\Gamma_1(N)$ of weight k and character χ_γ .

Proposition 3.5

Let $a/c \in \mathbb{Q}$ with $(a, c) = 1$ be a cusp of $\Gamma_1(N)$ and let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$.

If a/c is regular then

$$f|_M(\tau) = \sum a_n q_{mt}^n$$

where n is integral and $n = a^2 j m t / N \pmod{m}$.

Suppose a/c is irregular. If km is even then

$$f|_M(\tau) = \sum a_n q_m^n$$

where n is integral and $n = km/2 - jm/4 \pmod{m}$. If km is odd then

$$f|_M(\tau) = \sum a_n q_2^n$$

with $n = 1 \pmod{2}$.

Proof: Let a/c be regular. Then

$$f|_M(T^t \tau) = f|_{MT^t}(\tau) = f|_{T_{a/c}M}(\tau) = \chi_\gamma(T_{a/c})f|_M(\tau).$$

Since $\chi_\gamma(T_{a/c})$ has order m it follows that $f|_M$ has an expansion in integral powers of q_{mt} . Write $f|_M(\tau) = \sum a_n q_{mt}^n$. Then $f|_M(T^t \tau) = \sum a_n e(nt/mt) q_{mt}^n$ so that $f|_M(T^t \tau) = \chi_\gamma(T_{a/c})f|_M(\tau)$ holds if and only if $e(n/m) = \chi_\gamma(T_{a/c})$ for all n with $a_n \neq 0$. But this is equivalent to $n = a^2 j m t / N \pmod{m}$.

If a/c is irregular then

$$f|_M(T\tau) = f|_{T_{a/c}M}(\tau) = \chi_\gamma(-T_{a/c})f|_{-M}(\tau) = (-1)^k \chi_\gamma(-T_{a/c})f|_M(\tau)$$

and the statement follows by a similar argument. \square

Proposition 3.6

Let $a/c \in \mathbb{Q}$ with $(a, c) = 1$ be a cusp of $\Gamma_1(N)$ and $\mu \in a\gamma + D^{c*}$.

If a/c is regular then $\mu^2/2 = -n/mt \pmod{1}$ where n is an integer such that $n = a^2 jmt/N \pmod{m}$.

Suppose a/c is irregular and $(-1)^k e(\text{sign}(D)/4) e(\gamma\mu) = 1$. If km is even then $\mu^2/2 = k/2 + j/4 \pmod{1}$. If km is odd then $\mu^2/2 = 1/2 \pmod{1}$.

Proof: Choose a Jordan decomposition of D and let x_c be the canonical coset representative of D^{c*} . Write $\mu = a\gamma + (x_c + c\beta)$. Then

$$\mu^2/2 = a^2\gamma^2/2 + x_c^2/2 + c^2\beta^2/2 + ac\gamma\beta + ax_c\gamma \pmod{1}.$$

Let a/c be regular. Then $tx_c^2/2 = tc^2\beta^2/2 = tac\gamma\beta = tax_c\gamma = 0 \pmod{1}$ and $mta^2\gamma^2/2 = 0 \pmod{1}$. Hence $mt\mu^2/2 = 0 \pmod{1}$. Define $n = -mt\mu^2/2 \pmod{m}$. Then $n = -mta^2\gamma^2/2 = a^2 jmt/N \pmod{m}$.

Now let a/c be irregular. Then $N = 4$, $a = 1 \pmod{2}$ and $c = 2 \pmod{4}$. The Jordan decomposition of D is of the form $2_{II}^{\epsilon_2 n_2} 4_{II}^{\epsilon_4 n_4}$ or $2_{t_2}^{\epsilon_2 n_2} 4_{II}^{\epsilon_4 n_4}$. In the first case $\text{sign}(D) = \text{oddtity}(D) = 0 \pmod{4}$ and $x_c = 0$ and in the second case $\text{sign}(D) = \text{oddtity}(D) = t_2 \pmod{4}$ and $x_c^2/2 = t_2/4 = \text{sign}(D)/4 \pmod{1}$. The formula for the norm of μ simplifies to

$$\mu^2/2 = \gamma^2/2 + x_c^2/2 + c\gamma\beta + x_c\gamma \pmod{1}.$$

Using $\gamma\mu = c\gamma^2/2 + x_c\gamma + c\gamma\beta \pmod{1}$ we obtain

$$\mu^2/2 = -\gamma^2/2 + x_c^2/2 + \gamma\mu \pmod{1}.$$

Finally $k/2 + \text{sign}(D)/4 + \gamma\mu = k/2 + x_c^2/2 + \gamma\mu = 0 \pmod{1}$ so that

$$\mu^2/2 = k/2 - \gamma^2/2 \pmod{1}.$$

This implies the last statement. □

Let F be the lift of f on γ . Then as above

$$F = \sum_{s \in \Gamma_1(N) \setminus P} F_s$$

with

$$F_s = \sum_{\substack{M \in \Gamma_1(N) \setminus \Gamma \\ M\infty = s}} f|_M \rho_D(M^{-1}) e^\gamma.$$

The functions F_s are given in the following theorem.

Theorem 3.7

Let $a/c \in \mathbb{Q}$ with $(a, c) = 1$ be a representative of s and $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$.

If s is regular write $f|_M = g_{mt,0} + \dots + g_{mt,mt-1}$ with $g_{mt,j}|_T = e(j/mt)g_{mt,j}$ and for $\mu \in a\gamma + D^{c*}$ define j_μ by $\mu^2/2 = -j_\mu/mt \pmod{1}$. Then

$$F_s = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\mu \in a\gamma + D^{c*}} e(d(\mu - a\gamma)_c^2/2) e(b\mu\gamma) e(-ab\gamma^2/2) \\ t g_{mt,j_\mu} \{ e^\mu + (-1)^k e(\text{sign}(D)/4) e^{-\mu} \}.$$

if $N > 2$ and

$$F_s = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\mu \in a\gamma + D^{c*}} e(d(\mu - a\gamma)_c^2/2) e(b\gamma\mu) e(-ab\gamma^2/2) \\ t g_{mt, j_\mu} e^\mu.$$

if $N = 2$.

If a/c is irregular then

$$F_s = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\mu \in a\gamma + D^{c*}} e(d(\mu - a\gamma)_c^2/2) e(b\mu\gamma) e(-ab\gamma^2/2) \\ \{1 + (-1)^k e(\text{sign}(D)/4) e(\mu\gamma)\} f|_M e^\mu.$$

Proof: Let s be regular and $N > 2$. The cosets of $\Gamma_1(N)$ sending ∞ to s are represented by $\pm MT^n$ where n ranges over a complete set of residues modulo $t = N/(c, N)$. Using $f|_{-1} = (-1)^k f$ and $\rho_D(-1)e^\gamma = e(\text{sign}(D)/4)e^{-\gamma}$ we get

$$F_s = \sum_{n \bmod t} f|_{MT^n} \rho_D(T^{-n}) \rho_D(M^{-1}) e^\gamma \\ + (-1)^k e(\text{sign}(D)/4) \sum_{n \bmod t} f|_{MT^n} \rho_D(T^{-n}) \rho_D(M^{-1}) e^{-\gamma}.$$

Now

$$\rho_D(M^{-1}) e^\gamma = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in D^{c*}} e(d\beta_c^2/2) e(b\beta\gamma) e(ab\gamma^2/2) e^{\alpha\gamma + \beta}$$

so that

$$F_s = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{n \bmod t} f|_{MT^n} \rho_D(T^{-n}) \\ \left\{ \sum_{\beta \in D^{c*}} e(d\beta_c^2/2) e(b\beta\gamma) e(ab\gamma^2/2) e^{\alpha\gamma + \beta} \right. \\ \left. + (-1)^k e(\text{sign}(D)/4) \sum_{\beta \in D^{c*}} e(d\beta_c^2/2) e(-b\beta\gamma) e(ab\gamma^2/2) e^{-\alpha\gamma + \beta} \right\}.$$

Replacing β by $-\beta$ in the last sum and putting $\mu = a\gamma + \beta$ gives

$$F_s = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\mu \in a\gamma + D^{c*}} \sum_{n \bmod t} f|_{MT^n} e(n\mu^2/2) \\ e(d(\mu - a\gamma)_c^2/2) e(b\mu\gamma) e(-ab\gamma^2/2) \{e^\mu + (-1)^k e(\text{sign}(D)/4) e^{-\mu}\}.$$

The formula now follows from

$$\sum_{n \bmod t} f|_{MT^n} e(n\mu^2/2) = t g_{mt, j_\mu}.$$

The case $N = 2$ is now clear.

Suppose s is irregular, i.e. $N = 4$, $a = 1 \pmod{2}$ and $c = 2 \pmod{4}$. We argue slightly differently here because $a\gamma + D^{c*} = -a\gamma - D^{c*}$ in this case. We have

$$\begin{aligned} F_s &= f|_M \rho_D(M^{-1})e^\gamma + f|_{-M} \rho_D(-M^{-1})e^{-\gamma} \\ &= f|_M \rho_D(M^{-1})e^\gamma + (-1)^k e(\text{sign}(D)/4) f|_M \rho_D(M^{-1})e^{-\gamma} \end{aligned}$$

and

$$\begin{aligned} &\rho_D(M^{-1})e^{-\gamma} \\ &= \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in D^{c*}} e(d\beta_c^2/2)e(-b\beta\gamma)e(ab\gamma^2/2)e^{-a\gamma+\beta} \\ &= \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in 2a\gamma + D^{c*}} e(d\beta_c^2/2)e(-b\beta\gamma)e(ab\gamma^2/2)e^{-a\gamma+\beta} \\ &= \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in D^{c*}} e(d(2a\gamma + \beta)_c^2/2)e(-b(2a\gamma + \beta)\gamma)e(ab\gamma^2/2)e^{a\gamma+\beta} \\ &= \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in D^{c*}} e(d\beta_c^2/2)e(c\gamma^2/2 + \gamma\beta)e(b\gamma\beta)e(ab\gamma^2/2)e^{a\gamma+\beta} \end{aligned}$$

because $d(2a\gamma + \beta)_c^2/2 = d\beta_c^2/2 + c\gamma^2/2 + \gamma\beta \pmod{1}$ and $-b(2a\gamma + \beta)\gamma = b\beta\gamma \pmod{1}$. Hence

$$\begin{aligned} F_s &= \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in D^{c*}} e(d\beta_c^2/2)e(b\gamma\beta)e(ab\gamma^2/2) \\ &\quad \{1 + (-1)^k e(\text{sign}(D)/4)e(c\gamma^2/2 + \gamma\beta)\} f|_M e^{a\gamma+\beta} \end{aligned}$$

Let $\mu = a\gamma + \beta$. Then $ab\gamma^2/2 + b\gamma\beta = -ab\gamma^2/2 + b\gamma\mu \pmod{1}$ and $c\gamma^2/2 + \gamma\beta = \gamma\mu \pmod{1}$ so that

$$\begin{aligned} F_s &= \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\mu \in a\gamma + D^{c*}} e(d(\mu - a\gamma)_c^2/2)e(b\mu\gamma)e(-ab\gamma^2/2) \\ &\quad \{1 + (-1)^k e(\text{sign}(D)/4)e(\mu\gamma)\} f|_M e^\mu. \end{aligned}$$

This proves the theorem. \square

The group $\Gamma(N)$ has index $N^3 \prod_{p|N} (1 - 1/p^2)$ in Γ and

$$\begin{cases} 3 & \text{if } N = 2, \\ (N^2/2) \prod_{p|N} (1 - 1/p^2) & \text{if } N > 2 \end{cases}$$

classes of cusps. Two cusps a/c and a'/c' in \mathbb{Q} with $(a, c) = (a', c') = 1$ are equivalent modulo $\Gamma(N)$ if and only if

$$\begin{pmatrix} a' \\ c' \end{pmatrix} = \pm \begin{pmatrix} a \\ c \end{pmatrix} \pmod{N}.$$

The width of a/c is N and the stabilizer of a/c in $\Gamma(N)$ is generated by $T_{a/c}$ defined as above.

Let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Then the cosets of $\Gamma(N) \backslash \Gamma$ sending ∞ to a/c are represented by

$$\begin{aligned} MT^n & \quad \text{if } N = 2, \\ \pm MT^n & \quad \text{if } N > 2 \end{aligned}$$

where n ranges over a complete set of residues modulo N .

Let D be a discriminant form of even signature and level dividing N and f a modular form on $\Gamma(N)$ of weight k .

The function $f|_M$ has a Fourier expansion in integral powers of q_N for all $M \in \Gamma$.

Let F be the lift of f on $\gamma \in D$. Then $F = \sum_{s \in \Gamma(N) \backslash P} F_s$ and as above we find

Theorem 3.8

Let $a/c \in \mathbb{Q}$ with $(a, c) = 1$ be a representative of s and $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Decompose $f|_M$ into T -eigenfunctions $g_{N,j}$ and for $\mu \in a\gamma + D^{c*}$ define j_μ by $\mu^2/2 = -j_\mu/N \pmod{1}$. Then

$$\begin{aligned} F_s = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\mu \in a\gamma + D^{c*}} e(d(\mu - a\gamma)_c^2/2) e(b\mu\gamma) e(-ab\gamma^2/2) \\ N g_{N,j_\mu} \{ e^\mu + (-1)^k e(\text{sign}(D)/4) e^{-\mu} \}. \end{aligned}$$

if $N > 2$ and

$$\begin{aligned} F_s = \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\mu \in a\gamma + D^{c*}} e(d(\mu - a\gamma)_c^2/2) e(b\gamma\mu) e(-ab\gamma^2/2) \\ N g_{N,j_\mu} e^\mu. \end{aligned}$$

if $N = 2$.

4 Induction from isotropic subgroups

Let L be an even lattice and M a sublattice of L of finite index. We show that a modular form on the discriminant form of L induces in a canonical way a modular form on the discriminant form of M .

Let D be a discriminant form of even signature. Let H be an isotropic subgroup of D and H^\perp the orthogonal complement of H in D . Then $D_H =$

H^\perp/H is a discriminant form of the same signature as D with $|D_H| = |D|/|H|^2$. Let $F_{D_H} = \sum_{\gamma \in D_H} F_{D_H, \gamma} e^\gamma$ be a modular form for ρ_{D_H} . Define

$$F = \sum_{\gamma \in H^\perp} F_{D_H, \gamma+H} e^\gamma.$$

Then

Theorem 4.1

F is a modular form for ρ_D .

Proof: It is sufficient to show that $F = \sum_{\gamma \in D} F_\gamma e^\gamma$ transforms correctly under the generators of Γ . This is clear for T .

For $\gamma \in H^\perp$ we have

$$\begin{aligned} F_\gamma|_S(\tau) &= F_{D_H, \gamma+H}|_S(\tau) \\ &= \frac{e(\text{sign}(D_H)/8)}{\sqrt{|D_H|}} \sum_{\beta \in D_H} e((\gamma+H)\beta) F_{D_H, \beta}(\tau) \\ &= \frac{e(\text{sign}(D_H)/8)}{|H|\sqrt{|D_H|}} |H| \sum_{\beta \in D_H} e((\gamma+H)\beta) F_{D_H, \beta}(\tau) \\ &= \frac{e(\text{sign}(D_H)/8)}{|H|\sqrt{|D_H|}} \sum_{\beta \in H^\perp} e(\gamma\beta) F_\beta(\tau) \\ &= \frac{e(\text{sign}(D)/8)}{\sqrt{|D|}} \sum_{\beta \in D} e(\gamma\beta) F_\beta(\tau). \end{aligned}$$

If $\gamma \notin H^\perp$ then $F_\gamma|_S(\tau) = 0$ and

$$\begin{aligned} \sum_{\beta \in D} e(\gamma\beta) F_\beta(\tau) &= \sum_{\beta \in H^\perp} e(\gamma\beta) F_\beta(\tau) \\ &= \sum_{\beta \in H^\perp/H} \sum_{\mu \in H} e(\gamma(\beta+\mu)) F_{\beta+\mu}(\tau) \\ &= \sum_{\beta \in D_H} e((\gamma+H)\beta) F_{D_H, \beta}(\tau) \sum_{\mu \in H} e(\gamma\mu) \\ &= 0 \end{aligned}$$

because $\sum_{\mu \in H} e(\gamma\mu)$ is the sum over a nontrivial character of H and therefore is 0. \square

We can easily construct some automorphisms of F . An automorphism of D stabilizes H as a set if and only if it stabilizes H^\perp . Such automorphisms act on $H^\perp/H = D_H$ so that we get a natural map from the stabilizer of H in $O(D)$ to $O(D_H)$. The inverse image of $O(F_{D_H})$ under this map is a subgroup of $O(F)$.

A standard situation in which the above result can be applied is the following. Let L be an even lattice of even signature and let M be a sublattice of L of

finite index. Then we have embeddings $M \subset L \subset L' \subset M'$ and $H = L/M$ is an isotropic subgroup of $D_M = M'/M$. The orthogonal complement of H in D_M is $H^\perp = L'/M$ and H^\perp/H is naturally isomorphic to $D_L = L'/L$. The theorem then shows that a modular form for the discriminant form of L induces a modular form for the discriminant form of the sublattice M .

5 Discriminant forms of squarefree level

Let D be a discriminant form of even signature and level N . Let f be a modular form on $\Gamma_0(N)$ of character χ_D . Then $F_{\Gamma_0(N),f,0}$ is a modular form for ρ_D which is invariant under $O(D)$. We show now that every modular form for ρ_D which is invariant under $O(D)$ can be obtained in this way if N is squarefree.

First we describe some properties of discriminant forms of prime level.

Proposition 5.1

Let D be a discriminant form of prime level and let β and γ be two nonzero elements in D of the same norm. Then β and γ are conjugate under $O(D)$.

Proof: Let p be the level of D . If p is an odd prime we can consider D as a vector space over \mathbb{F}_p with a quadratic form. The statement then follows from Witt's theorem (cf. [O]). For $p = 2$ we can argue as follows. The statement is true if D has 2-rank 2. Suppose that the 2-rank is ≥ 4 . If $\beta^2/2 = 0 \pmod{1}$ then there is an $\alpha \in D$ such that $\alpha\beta = 1/2 \pmod{1}$. Replacing α by $\alpha + \beta$ if necessary we can assume that $\alpha^2/2 = 0 \pmod{1}$. Similarly if $\beta^2/2 = 1/2 \pmod{1}$ we can find $\alpha \in D$ such that $\alpha\beta = \alpha^2/2 = 1/2 \pmod{1}$. In both cases D decomposes into the orthogonal sum $D = \langle \alpha, \beta \rangle \oplus \langle \alpha, \beta \rangle^\perp$. This proves the proposition for $p = 2$. \square

For a discriminant form D we denote by I the subset of isotropic elements and by I_k the isotropic elements of order k .

Proposition 5.2

Let D be a discriminant form of prime level p . Then we have for $\gamma \in I_p$

$$\sum_{\beta \in I_p} e(\beta\gamma) = |I_p| - \frac{|D|}{p}.$$

Proof: We prove the statement in the case that D has Jordan decomposition $p^{\epsilon n}$ with p odd and $n \geq 3$. The other cases are similar. We can assume that

$$D = D' \oplus D''$$

where D' is generated by γ_1, γ_2 with $\gamma_1^2/2 = 1/p \pmod{1}$, $\gamma_2^2/2 = -1/p \pmod{1}$, $\gamma_1\gamma_2 = 0$, i.e. has Jordan decomposition $p^{\epsilon' 2}$ with $\epsilon' = \begin{pmatrix} 2 \\ p \end{pmatrix} \begin{pmatrix} -2 \\ p \end{pmatrix} = \begin{pmatrix} -1 \\ p \end{pmatrix}$, and contains γ , and D'' has Jordan decomposition $p^{\epsilon''(n-2)}$ with $\epsilon'' = \begin{pmatrix} -1 \\ p \end{pmatrix} \epsilon$. Then

$$\gamma^\perp = \langle \gamma \rangle \oplus D''$$

so that by Proposition 3.2 in [S1]

$$\begin{aligned} |\gamma^\perp \cap I| &= p \begin{cases} p^{n-3} + \epsilon'' \left(\frac{-1}{p}\right)^{(n-2)/2} (p^{(n-2)/2} - p^{(n-4)/2}) & \text{if } n \text{ is even} \\ p^{n-3} & \text{if } n \text{ is odd} \end{cases} \\ &= \begin{cases} p^{n-2} + \epsilon \left(\frac{-1}{p}\right)^{n/2} (p^{n/2} - p^{(n-2)/2}) & \text{if } n \text{ is even} \\ p^{n-2} & \text{if } n \text{ is odd} \end{cases} \end{aligned}$$

Define $A_j = \{\beta \in I \mid \gamma\beta = j/p \pmod{1}\}$. Then

$$I = \bigcup_{j=0}^{p-1} A_j.$$

We have $A_0 = \gamma^\perp \cap I$ and $|A_1| = \dots = |A_{p-1}| = m$ because multiplication by elements in $(\mathbb{Z}/p\mathbb{Z})^*$ maps the sets into each other. Hence applying again Proposition 3.2 in [S1] we obtain

$$m = \frac{|I| - |A_0|}{p-1} = p^{n-2}.$$

It follows

$$\begin{aligned} \sum_{\beta \in I} e(\gamma\beta) &= |A_0| + m \sum_{j=1}^{p-1} e(j/p) \\ &= |A_0| - m \\ &= |A_0| - p^{n-2} \\ &= |I| - p^{n-1} \\ &= |I_p| + 1 - \frac{|D|}{p} \end{aligned}$$

This proves the statement. \square

Let D be a discriminant form of squarefree level N . Then Proposition 5.1 implies that two elements of D are in the same orbit under $O(D)$ if and only if they have the same norm and order. Let $F = \sum_{\gamma \in D} F_\gamma e^\gamma$ be a modular form for ρ_D which is invariant under $O(D)$. Then the components F_γ with isotropic γ are modular forms on $\Gamma_0(N)$ with character χ_D .

Proposition 5.3

Let D be a discriminant form of squarefree level N and $F = \sum_{\gamma \in D} F_\gamma e^\gamma$ a modular form for ρ_D which is invariant under $O(D)$. Then the complex vector space W spanned by the components F_γ , $\gamma \in D$ is generated by the functions $F_0|_M$, $M \in \Gamma$. In particular $F = 0$ if $F_0 = 0$.

Proof: Clearly W contains the functions $F_0|_M$, $M \in \Gamma$. Let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Then $F_0|_M$ decomposes into T -eigenfunctions, i.e.

$$F_0|_M = g_{t,0} + \dots + g_{t,t-1}$$

where $t = N/(c^2, N)$ is the width of a/c and $g_{t,j}|_T = e(j/t)g_{t,j}$. The function $g_{t,j}$ can be written as

$$g_{t,j} = \frac{1}{t} \sum_{n \bmod t} e(-jn/t) F_0|_{MT^n}$$

hence is in W . Now let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ with $c|N$ and $d = 0 \bmod c'$ where $c' = N/c$. Then

$$F_0|_M = \bar{\xi}(M^{-1}) \frac{1}{\sqrt{|D_{c'}|}} \sum_{\gamma \in D_{c'}} F_\gamma.$$

Since F_γ is determined by the norm and order of γ we see that F_γ can be written as a linear combination of functions $F_0|_M$, $M \in \Gamma$. \square

The main result of this section is the following.

Theorem 5.4

Let D be a discriminant form of squarefree level N and $F = \sum_{\gamma \in D} F_\gamma e^\gamma$ a modular form for ρ_D which is invariant under $O(D)$. Let N_R be the product over the primes with nonvanishing I_p . For $k|N_R$ define $F_k = F_\gamma$ where γ is any element in I_k . Then the functions F_k span the subspace W_0 of W with T -eigenvalue $e(0)$. Define

$$\begin{aligned} \Phi : W_0 &\longrightarrow W_0 \\ f &\longmapsto 0\text{-component of } F_{\Gamma_0(N),f,0} \end{aligned}$$

Then

$$\Phi(F_k) = \sum_{j|N_R} a_{jk} F_j$$

with

$$\begin{aligned} a_{jk} &= \frac{N}{|D|} |I_j| \sum_{c|(N/j, N/k)} \frac{|D_c|}{c} \\ &= \frac{N}{|D|} \prod_{p|j} |I_p| \prod_{p|(N/j, N/k)} \left(1 + \frac{|D_p|}{p}\right). \end{aligned}$$

The matrix $A = (a_{jk})$ has determinant

$$\det(A) = \left(\frac{N}{|D|}\right)^{\sigma(N_R)} \left(\sum_{d|N/N_R} \frac{|D_d|}{d}\right)^{\sigma(N_R)} \prod_{d|N_R} |I_d| \prod_{d|N_R} \frac{|D_d|}{d}.$$

In particular Φ is invertible.

Proof: Let $\gamma \in I_k$ where $k|N_R$. Then

$$\Phi(F_\gamma) = \sum_{s \in \Gamma_0(N) \setminus P} \Phi_s(F_\gamma)$$

with

$$\Phi_s(F_\gamma) = \sum_{\substack{M \in \Gamma_0(N) \setminus \Gamma \\ M\infty=s}} F_\gamma|_M \langle \rho_D(M^{-1})e^0, e^0 \rangle.$$

The cusps of $\Gamma_0(N)$ are given by $1/c$, where c ranges over the positive divisors of N , and the cusp $1/c$ has width $c' = N/c$.

Let $c|N$. Choose $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ with $d \equiv 1 \pmod{c}$ and $d \equiv 0 \pmod{c'}$. Then

$$\begin{aligned} \Phi_{1/c}(F_\gamma) &= \sum_{n \pmod{c'}} F_\gamma|_{MT^n} \langle \rho_D((MT^n)^{-1})e^0, e^0 \rangle \\ &= \sum_{n \pmod{c'}} \sum_{\alpha \in D} F_\alpha \langle \rho_D(MT^n)e^\alpha, e^\gamma \rangle \langle \rho_D((MT^n)^{-1})e^0, e^0 \rangle \end{aligned}$$

$\Phi_{1/c}(F_\gamma)$ is T -invariant so that the contributions of the $\alpha \in D$ with nonzero norm cancel each other. Hence

$$\Phi_{1/c}(F_\gamma) = \frac{N}{c} \sum_{\substack{\alpha \in D \\ \alpha^2/2 \equiv 0 \pmod{1}}} F_\alpha \langle e^\alpha, \rho_D(M^{-1})e^\gamma \rangle \langle \rho_D(M^{-1})e^0, e^0 \rangle.$$

Now

$$\begin{aligned} \rho_D(M^{-1})e^\gamma &= \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in D_{c'}} e(b\beta\gamma) e^{a\gamma+\beta} \\ &= \xi(M^{-1}) \frac{\sqrt{|D_c|}}{\sqrt{|D|}} \sum_{\beta \in D_{c'}} e(-c^{-1}\beta\gamma) e^{a\gamma+\beta} \end{aligned}$$

where c^{-1} is the inverse of c modulo c' so that

$$\begin{aligned} \Phi_{1/c}(F_\gamma) &= \frac{N}{c} \frac{|D_c|}{|D|} \sum_{\substack{\alpha \in D \\ \alpha^2/2 \equiv 0 \pmod{1}}} F_\alpha \sum_{\beta \in D_{c'}} e(c^{-1}\beta\gamma) \langle e^\alpha, e^{a\gamma+\beta} \rangle \\ &= \frac{N}{c} \frac{|D_c|}{|D|} \sum_{\substack{\alpha \in (\gamma+D_{c'}) \\ \alpha^2/2 \equiv 0 \pmod{1}}} e(c^{-1}\alpha\gamma) F_\alpha \end{aligned}$$

and

$$\Phi(F_\gamma) = \sum_{c|N} \frac{N}{c} \frac{|D_c|}{|D|} \sum_{\substack{\alpha \in (\gamma+D_{c'}) \\ \alpha^2/2 \equiv 0 \pmod{1}}} e(c^{-1}\alpha\gamma) F_\alpha.$$

Since

$$\sum_{\substack{\alpha \in (\gamma + D_{c'}) \\ \alpha^2/2 \equiv 0 \pmod{1}}} e(c^{-1}\alpha\gamma)F_\alpha = \sum_{j|N_R} F_j \sum_{\alpha \in (\gamma + D_{c'}) \cap I_j} e(c^{-1}\alpha\gamma)$$

we get

$$\Phi(F_k) = \sum_{j|N_R} a_{jk}F_j$$

with

$$a_{jk} = \sum_{c|N} \frac{N}{c} \frac{|D_c|}{|D|} \sum_{\alpha \in (\gamma + D_{c'}) \cap I_j} e(c^{-1}\alpha\gamma).$$

The elements in $\gamma + D_{c'}$ have order $(k, c)d$ where $d|c'$ and the elements in I_j have order j . Hence

$$\begin{aligned} a_{jk} &= \sum_{\substack{c|N \\ (k,c)|j|(k,c)c'}} \frac{N}{c} \frac{|D_c|}{|D|} \sum_{\alpha \in (\gamma + D_{c'}) \cap I_j} e(c^{-1}\alpha\gamma) \\ &= \sum_{\substack{c|N \\ (k,c)|j|(k,c)c'}} \frac{N}{c} \frac{|D_c|}{|D|} \sum_{\alpha \in I_{j/(k,c)}} e(c^{-1}\alpha\gamma) \\ &= \sum_{\substack{c|N \\ (j,c)=(k,c)}} \frac{N}{c} \frac{|D_c|}{|D|} \sum_{\alpha \in I_{(j,c')}} e(\alpha\gamma). \end{aligned}$$

We can evaluate the last sum using Proposition 5.2. We get

$$a_{jk} = \frac{N}{|D|} \sum_{\substack{c|N \\ (j,c)=(k,c)}} \frac{|D_c|}{c} \prod_{\substack{p|(j,c') \\ p \nmid k}} |I_p| \prod_{\substack{p|(j,c') \\ p \nmid k}} \left(|I_p| - \frac{|D_p|}{p} \right).$$

Induction on the number of prime divisors of N shows

$$\begin{aligned} a_{jk} &= \frac{N}{|D|} \prod_{p|j} |I_p| \sum_{c|(N/j, N/k)} \frac{|D_c|}{c} \\ &= \frac{N}{|D|} |I_j| \prod_{p|(N/j, N/k)} \left(1 + \frac{|D_p|}{p} \right) \\ &= \frac{N}{|D|} \prod_{p|N/N_R} \left(1 + \frac{|D_p|}{p} \right) |I_j| \prod_{p|(N_R/j, N_R/k)} \left(1 + \frac{|D_p|}{p} \right). \end{aligned}$$

We can write $A = (a_{jk})$ as

$$A = \frac{N}{|D|} \prod_{p|N/N_R} \left(1 + \frac{|D_p|}{p} \right) BC$$

where $B = (b_{jk})$ is a diagonal matrix with entries

$$b_{jk} = \delta_{jk} |I_j|$$

and $C = (c_{jk})$ is a symmetric matrix with entries

$$c_{jk} = \prod_{p|(N_R/j, N_R/k)} \left(1 + \frac{|D_p|}{p} \right).$$

We have

$$\det(B) = \prod_{d|N_R} |I_d|$$

and

$$\det(C) = \prod_{d|N_R} \frac{|D_d|}{d}$$

so that

$$\det(A) = \left(\frac{N}{|D|} \right)^{\sigma(N_R)} \left(\sum_{d|N/N_R} \frac{|D_d|}{d} \right)^{\sigma(N_R)} \prod_{d|N_R} |I_d| \prod_{d|N_R} \frac{|D_d|}{d}$$

where $\sigma(N_R)$ denotes the number of divisors of N_R . This finishes the proof of the theorem. \square

Proposition 5.3 and Theorem 5.4 imply

Corollary 5.5

Let D be a discriminant form of squarefree level N and F a modular form for ρ_D which is invariant under $O(D)$. Then $F = F_{\Gamma_0(N), f, 0}$ for a suitable modular form f on $\Gamma_0(N)$ with character χ_D .

The last result is also stated as Theorem 4.2.17 in [Ba]. However the proof given there is incomplete as the author does not show that the matrix A is invertible.

6 Some new generalized Kac-Moody algebras

In this section we use the above results to construct some modular forms for the Weil representation with nonnegative integral coefficients and reflective poles. The theta lifts of these modular forms are holomorphic automorphic products of singular weight. They give new examples of generalized Kac-Moody algebras with automorphic denominator identity.

The automorphic forms of this section are related to automorphisms of the Leech lattice with cycle shapes 2^{12} , 3^8 and 3.21 . However we will see that the first two examples can be constructed more naturally from the Niemeier lattices with root system D_{12}^2 and E_8^3 .

We have used the computer algebra systems Magma [BCP] and PARI/GP [P] in the proof of some statements.

We begin with the case of cycle shape 3^8 because this is the simplest example.

Cycle shape 3^8

Let Λ be the Leech lattice and g an automorphism of Λ of cycle shape 3^8 . Then the fixed point sublattice Λ^g of g is a primitive sublattice of Λ isomorphic to $\sqrt{3}E_8$.

The lattice $\sqrt{3}E_8$ has genus $II_{8,0}(3^{+8})$ and represents the unique class in this genus. The orthogonal group of $\sqrt{3}E_8$ acts transitively on the vectors of norm 6, 12 and 18. The vectors of norm 24 decompose into 2 orbits of length 240 and 17280. It follows that $O(\sqrt{3}E_8)$ has 5 orbits on the discriminant form of $\sqrt{3}E_8$ which we describe in the following table.

| norm | length | order | name |
|------|--------|-------|-------|
| 0 | 1 | 1 | 0_0 |
| 0 | 2240 | 3 | 0 |
| 1/3 | 240 | 3 | 1_S |
| | 1920 | 3 | 1_L |
| 2/3 | 2160 | 3 | 2 |

The centralizer $C(g)$ of g in $O(\Lambda)$ does not induce the full orthogonal group of Λ^g . It acts transitively on the vectors of norm 6, 12 and 18 in Λ^g and the vectors of norm 24 decompose into 3 orbits of length 240, 2160 and 15120. The orbit 1_L in the above table splits into 2 orbits under $C(g)$ which we denote by 1_{L_S} and 1_{L_L} . They contain 240 respectively 1680 elements.

The orthogonal complement $\Lambda^{g\perp}$ of Λ^g in Λ is isomorphic to $A_2 \otimes E_8$.

The lattice $A_2 \otimes E_8$ is 3-modular and has genus $II_{16,0}(3^{+8})$. The orthogonal group of $A_2 \otimes E_8$ has 5 orbits on the discriminant form of $A_2 \otimes E_8$ which are as follows.

| norm | length | order | name |
|------|--------|-------|-------|
| 0 | 1 | 1 | 0_0 |
| 0 | 2240 | 3 | 0 |
| 1/3 | 2160 | 3 | 1 |
| 2/3 | 240 | 3 | 2_S |
| | 1920 | 3 | 2_L |

The theta functions of the corresponding cosets are given by

$$\begin{aligned}
 \theta_{0_0}(\tau) &= 1 + 720q^2 + 13440q^3 + 97200q^4 + 455040q^5 + 1714320q^6 + \dots \\
 \theta_0(\tau) &= 6q + 765q^2 + 12960q^3 + 97863q^4 + 463068q^5 + 1672083q^6 + \dots \\
 \theta_1(\tau) &= 45q^{4/3} + 2232q^{7/3} + 27306q^{10/3} + 170064q^{13/3} + \dots \\
 \theta_{2_S}(\tau) &= 3q^{2/3} + 168q^{5/3} + 6009q^{8/3} + 51960q^{11/3} + 288600q^{14/3} + \dots \\
 \theta_{2_L}(\tau) &= 216q^{5/3} + 5697q^{8/3} + 52920q^{11/3} + 287820q^{14/3} + \dots
 \end{aligned}$$

Under the action of the centralizer $C(g)$ the orbit 2_L splits into 2 orbits of length 240 and 1680 which we denote by 2_{L_S} and 2_{L_L} .

Since Λ^g is a primitive sublattice of Λ we have a natural isomorphism of groups

$$i : D_{\Lambda^g} \rightarrow D_{\Lambda^{g\perp}}$$

satisfying $i(\gamma)^2/2 = -\gamma^2/2$ for all $\gamma \in D_{\Lambda^g}$. The map i sends the orbit 1_S into 2_{L_S} and 1_{L_S} into 2_S because Λ has no vectors of norm 2.

Let H be a lattice with Gram matrix $\begin{pmatrix} -2 & 3 \\ 3 & 0 \end{pmatrix}$. Then H has genus $II_{1,1}(9^{-1})$ and we can consider H as a sublattice of $II_{1,1}$ of index 3. Hence $\theta_{\Lambda^{g\perp}}/3\Delta$ induces a modular form $F_{\theta_{\Lambda^{g\perp}}/3\Delta}$ of weight -4 for the Weil representation of $L = \Lambda^g \oplus H$ (cf. the remarks at the end of Section 2, Theorem 4.1 and the remark at the end of Section 4). We decompose $D_L = D_{\Lambda^g} \oplus D_H$. Then the components of $F_{\theta_{\Lambda^{g\perp}}/3\Delta}$ are given by

$$F_{\theta_{\Lambda^{g\perp}}/3\Delta, \gamma} = \theta_{i(\gamma_{\Lambda^g}) + \Lambda^{g\perp}}/3\Delta$$

if $\gamma = (\gamma_{\Lambda^g}, \gamma_H)$ with $\gamma_H \in D_H^3$ and $F_{\theta_{\Lambda^{g\perp}}/3\Delta, \gamma} = 0$ otherwise.

The function $\eta_g(\tau) = \eta(3\tau)^8$ is a modular form for $\Gamma_0(9)$ of weight 4 with trivial character. This function is related to the theta functions by

$$(\theta_{2_S} - \theta_{2_L})/3\Delta(\tau) = 1/\eta_g(\tau/3).$$

We decompose

$$1/\eta_g(\tau/9) = g_0(\tau) + \dots + g_8(\tau)$$

where $g_j|_T(\tau) = e(j/9)g_j(\tau)$. Note that the only nonvanishing g_j are g_2, g_5 and g_8 . Then the lifting of $1/\eta_g$ on the trivial subgroup of D_L is given by

$$F_{\Gamma_0(9), 1/\eta_g, 0} = \frac{3}{\eta_g} e^0 - \sum_{\gamma \in D_L^3} \frac{1}{\eta_g} e^\gamma + 3 \sum_{\gamma \in D_L} g_{j_\gamma} e^\gamma$$

where j_γ is defined by $j_\gamma/9 = -\gamma^2/2 \pmod{1}$ (cf. Theorem 3.2). Let

$$F = F_{\theta_{\Lambda^{g\perp}}/3\Delta} + \frac{1}{3} F_{\Gamma_0(9), 1/\eta_g, 0}.$$

Then we have

Theorem 6.1

F is a modular form for ρ_{D_L} of weight -4 . The components of F are given by

$$\begin{aligned} F_0 &= \frac{1}{\eta_g} + \frac{1}{3} \left(\frac{\theta_{0_0}}{\Delta} - \frac{1}{\eta_g} \right) \\ &= q^{-1} + 8 + 108q + 1072q^2 + 8790q^3 + 64512q^4 + 440176q^5 + \dots \end{aligned}$$

and

$$\begin{aligned} F_\gamma &= \frac{1}{3} \left(\frac{\theta_{0_0}}{\Delta} - \frac{1}{\eta_g} \right) \\ &= 8 + 108q + 1064q^2 + 8790q^3 + 64512q^4 + 440132q^5 + \dots \end{aligned}$$

if $\gamma \in D_L^3 \setminus \{0\}$,

$$F_\gamma = \frac{\theta_0}{3\Delta} = 2 + 303q + 11088q^2 + 225321q^3 + 3204240q^4 + \dots$$

if $\gamma^2/2 = 0 \pmod{1}$ and $\gamma \notin D_L^3$,

$$F_\gamma = \frac{\theta_{2_S}}{3\Delta} = q^{-1/3} + 80q^{2/3} + 3671q^{5/3} + 86736q^{8/3} + 1365702q^{11/3} + \dots$$

if $\gamma^2/2 = 1/3 \pmod{1}$ and $\gamma = (\gamma_{\Lambda^g}, \gamma_H)$ with $\gamma_{\Lambda^g} \in 1_{L_S}$,

$$F_\gamma = \frac{\theta_{2_L}}{3\Delta} = 72q^{2/3} + 3627q^{5/3} + 86544q^{8/3} + 1364976q^{11/3} + \dots$$

if $\gamma^2/2 = 1/3 \pmod{1}$ and $\gamma = (\gamma_{\Lambda^g}, \gamma_H)$ with $\gamma_{\Lambda^g} \in 1_S \cup 1_{L_L}$,

$$F_\gamma = \frac{\theta_1}{3\Delta} = 15q^{1/3} + 1104q^{4/3} + 31818q^{7/3} + 564192q^{10/3} + \dots$$

if $\gamma^2/2 = 2/3 \pmod{1}$,

$$F_\gamma = g_8 = q^{-1/9} + 192q^{8/9} + 7704q^{17/9} + 164560q^{26/9} + \dots$$

if $\gamma^2/2 = 1/9 \pmod{1}$,

$$F_\gamma = g_5 = 44q^{5/9} + 2464q^{14/9} + 62337q^{23/9} + 1020416q^{32/9} + \dots$$

if $\gamma^2/2 = 4/9 \pmod{1}$,

$$F_\gamma = g_2 = 8q^{2/9} + 726q^{11/9} + 22528q^{20/9} + 417140q^{29/9} + \dots$$

if $\gamma^2/2 = 7/9 \pmod{1}$.

The Fourier coefficients of the F_γ are nonnegative rational integers.

The poles of F define a reflection group of L which has a Weyl vector (cf. Theorems 12.1 and 10.4 in [B2]). For our purposes it is more convenient to consider the corresponding dual notions. We define W as the reflection group of the dual lattice L' generated by the roots

$$\begin{aligned} \alpha &\in L \text{ with } \alpha^2 = 2, \\ \alpha &\in L' \text{ with } \alpha^2 = 2/3 \text{ and } \alpha_{\Lambda^g} + \Lambda^g \in 1_{L_S}, \\ \alpha &\in L' \text{ with } \alpha^2 = 2/9. \end{aligned}$$

Then W has a Weyl vector ρ in $\mathbb{R} \otimes L$, i.e. the simple roots of W are the roots α satisfying $\rho\alpha = -\alpha^2/2$. The vector 3ρ is a primitive norm 0 vector in L' and $3\rho + L$ is in $D_L^3 \setminus \{0\}$. The primitive norm 0 vectors in H' are conjugate under $O(H')$. We can choose any of these vectors for 3ρ .

Let

$$M = L \oplus II_{1,1}.$$

Then F defines a modular form for ρ_{D_M} . The singular theta correspondence (cf. [B2], Theorem 13.3) maps F to an automorphic form Ψ on the Grassmannian of 2-dimensional negative definite subspaces of $\mathbb{R} \otimes M$. The lattice M has one orbit of primitive norm 0 vectors of level 1 under $O(M)$ and has no primitive norm 0 vectors of level 9.

Theorem 6.2

The automorphic form Ψ is holomorphic and has singular weight 4. The level 1 expansion is given by

$$e((\rho, Z)) \prod_{\alpha \in L'^+} (1 - e((\alpha, Z)))^{[F_{\alpha+L}](-\alpha^2/2)} = \sum_{w \in W} \det(w) \eta_g((w\rho, Z))$$

Proof: The level 1 product expansion of Ψ is

$$e((\rho, Z)) \prod_{\alpha \in L'^+} (1 - e((\alpha, Z)))^{[F_{\alpha+L}](-\alpha^2/2)}.$$

Since Ψ is holomorphic and has singular weight its Fourier expansion is supported only on norm 0 vectors. Furthermore Ψ is antisymmetric under the Weyl group W because the roots of W have multiplicity 1. It follows that Ψ has the sum expansion

$$\sum_{w \in W} \det(w) e((w\rho, Z)) \prod_{n > 0} (1 - e((3nw\rho, Z)))^8.$$

This proves the theorem. □

The above identity is also the denominator identity of a generalized Kac-Moody algebra whose real simple roots are the simple roots of W and imaginary simple roots are the positive integral multiples of 3ρ with multiplicity 8. The root lattice is L' and the multiplicity of a root α in L' is given by $[F_{\alpha+L}](-\alpha^2/2)$. This Lie algebra can also be constructed by orbifolding the fake monster algebra with a lift of g (cf. [B1], [S1]).

We describe a slight variation of the above construction which simplifies some aspects. Let N be the Niemeier lattice with root system E_8^3 (cf. [CS]). Then the direct product $O(E_8)^3$ is a normal subgroup of $O(N)$ with quotient S_3 . Let g be a permutation of the three E_8 -components of N of order 3. Then g has cycle shape 3^8 and fixed point sublattice N^g isomorphic to $\sqrt{3}E_8$. The orthogonal complement $N^{g\perp}$ is isomorphic to $A_2 \otimes E_8$. Here the centralizer $C(g)$ of g in $O(N)$ induces the full orthogonal group of N^g so that the natural isomorphism $i : D_{N^g} \rightarrow D_{N^{g\perp}}$ maps the orbits of the corresponding size into each other. If we construct F as above we can describe the components entirely in terms of the orbits of $O(\sqrt{3}E_8)$ on $D_{\sqrt{3}E_8}$. This also leads to a simpler description of W . For example the roots of norm $2/3$ then correspond to the elements in 1_S .

Cycle shape 2^{12}

Let g be an automorphism of Λ of cycle shape 2^{12} . Then Λ^g is a primitive sublattice of Λ isomorphic to $\sqrt{2}D_{12}^+$.

The lattice $\sqrt{2}D_{12}^+$ has genus $II_{12,0}(2_4^{+12})$ and there are 3 classes in this genus given by A_1^{12} , $\sqrt{2}E_8 \oplus A_1^4$ and $\sqrt{2}D_{12}^+$. The orthogonal group of $\sqrt{2}D_{12}^+$ acts transitively on the vectors of norm 4, 6 and 10. The vectors of length 8 decompose into 2 orbits of length 24 and 7920 and the vectors of length 12

decompose into 2 orbits of length 5280 and 59136. This implies that $O(\sqrt{2}D_{12}^+)$ has 7 orbits on the discriminant form of $\sqrt{2}D_{12}^+$. They are described in the following table.

| norm | length | order | name |
|------|--------|-------|-------|
| 0 | 1 | 1 | 0_0 |
| 0 | 1 | 2 | 0_S |
| | 990 | 2 | 0_L |
| 1/4 | 1024 | 2 | 1 |
| 1/2 | 132 | 2 | 2_S |
| | 924 | 2 | 2_L |
| 3/4 | 1024 | 2 | 3 |

The centralizer $C(g)$ of g in $O(\Lambda)$ does not induce the full orthogonal group of Λ^g . It acts transitively on the vectors of norm 4, 6 and 10 in Λ^g while the vectors of norm 8 decompose into 2 orbits of length 24 and 7920 and the vectors of norm 12 decompose into 3 orbits of length 5280, 8448 and 50688. Therefore the orbit 2_L in the above table splits into 2 orbits under $C(g)$ which we denote by 2_{L_S} and 2_{L_L} . They contain 132 respectively 792 elements.

The orthogonal complement $\Lambda^{g\perp}$ of Λ^g in Λ is Λ^{-g} and is also isomorphic to $\sqrt{2}D_{12}^+$. The theta functions of the cosets are given by

$$\begin{aligned}
\theta_{0_0}(\tau) &= 1 + 264q^2 + 2048q^3 + 7944q^4 + 24576q^5 + 64416q^6 + \dots \\
\theta_{0_S}(\tau) &= 24q + 3808q^3 + 50448q^5 + 268224q^7 + 947896q^9 + \dots \\
\theta_{0_L}(\tau) &= 8q + 256q^2 + 1952q^3 + 8192q^4 + 25008q^5 + 62464q^6 + \dots \\
\theta_1(\tau) &= 24q^{5/4} + 464q^{9/4} + 2904q^{13/4} + 11088q^{17/4} + 32032q^{21/4} + \dots \\
\theta_{2_S}(\tau) &= 2q^{2/4} + 40q^{6/4} + 876q^{10/4} + 4048q^{14/4} + 14650q^{18/4} + \dots \\
\theta_{2_L}(\tau) &= 64q^{6/4} + 768q^{10/4} + 4224q^{14/4} + 14848q^{18/4} + 40128q^{22/4} + \dots \\
\theta_3(\tau) &= 2q^{3/4} + 132q^{7/4} + 1254q^{11/4} + 5964q^{15/4} + 19338q^{19/4} + \dots
\end{aligned}$$

The lattice Λ^g is a primitive sublattice of Λ so that we have a natural isomorphism of groups

$$i : D_{\Lambda^g} \rightarrow D_{\Lambda^{g\perp}}$$

satisfying $i(\gamma)^2/2 = -\gamma^2/2$ for all $\gamma \in D_{\Lambda^g}$. Since Λ has no roots i maps the orbit 2_{L_S} of D_{Λ^g} into the orbit 2_S of $D_{\Lambda^{g\perp}}$ and the orbit 2_S into 2_{L_S} .

Let K be a lattice isomorphic to $\sqrt{2}D_{12}$. Then K has genus $II_{12,0}(2_{II}^{-10}4_{II}^{-2})$. There are 2 classes in this genus and the other class is given by $\sqrt{2}(E_8 \oplus D_4)$. The group $O(K)$ has 16 orbits on D_K . We describe them in the following tables.

| norm | length | order | norm | length | order |
|------|--------|-------|------|--------|-------|
| 0 | 1 | 1 | 1/4 | 24 | 4 |
| 0 | 1 | 2 | | 1584 | 4 |
| | 990 | 2 | | 4096 | 4 |
| | 990 | 2 | | 440 | 4 |
| | 2 | 2 | 3/4 | 440 | 4 |
| 1/2 | 132 | 2 | | 4096 | 4 |
| | 1848 | 2 | | 1548 | 4 |
| | 132 | 2 | | 24 | 4 |

Λ^g contains a unique sublattice isomorphic to K . This is the sublattice generated by the 264 vectors of norm 4 in Λ^g .

We consider K as a sublattice of Λ^g . Then $H = \Lambda^g/K$ is a subgroup of D_K^2 of order 2. Note that the elements in D_K of norm 0 or 1/2 all have order dividing 2 and therefore are in H^\perp . The function $\theta_{\Lambda^g/2\Delta}$ induces a modular form $F_{\theta_{\Lambda^g/2\Delta}}$ of weight -6 for the Weil representation of K . The components are given by

$$F_{\theta_{\Lambda^g/2\Delta}, \gamma} = \theta_{i(\gamma+H)+\Lambda^g/2\Delta}$$

if $\gamma \in H^\perp$ and $F_{\theta_{\Lambda^g/2\Delta}, \gamma} = 0$ otherwise.

The function $\eta_g(\tau) = \eta(2\tau)^{12}$ is a modular form for $\Gamma_0(4)$ of weight 6 with trivial character. We have the following relations

$$\begin{aligned} (\theta_{0_0} - \theta_{0_S})/\Delta(\tau) &= 1/\eta_g(\tau) \\ (\theta_{2_S} - \theta_{2_L})/2\Delta(\tau) &= 1/\eta_g(\tau/2) \\ (\theta_1 + \theta_3)/2\Delta(\tau) &= 1/\eta_g(\tau/4) \end{aligned}$$

We decompose $1/\eta_g(\tau/4)$ as

$$1/\eta_g(\tau/4) = g_0(\tau) + g_1(\tau) + g_2(\tau) + g_3(\tau)$$

with $g_j|_T(\tau) = e(j/4)g_j(\tau)$. Then the lifting of $1/\eta_g$ on the trivial subgroup of D_K is given by

$$F_{\Gamma_0(4), 1/\eta_g, 0} = \frac{2}{\eta_g} e^0 - \frac{1}{2} \sum_{\gamma \in D_K^2} \frac{1}{\eta_g} e^\gamma + 2 \sum_{\gamma \in D_K} g_{j_\gamma} e^\gamma$$

where j_γ is defined by $j_\gamma/4 = -\gamma^2/2 \pmod{1}$. The lift of $1/\eta_g$ on H is

$$F_{\Gamma_0(4), 1/\eta_g, H} = 2 \sum_{\gamma \in H} \frac{1}{\eta_g} e^\gamma - \sum_{\gamma \in D_K^2} \frac{1}{\eta_g} e^\gamma + 4 \sum_{\gamma \in H^\perp} g_{j_\gamma} e^\gamma.$$

Finally we define

$$F = F_{\theta_{\Lambda^g/2\Delta}} + \frac{1}{2} F_{\Gamma_0(4), 1/\eta_g, 0} - \frac{1}{4} F_{\Gamma_0(4), 1/\eta_g, H}.$$

Then the above relations imply

Theorem 6.3

F is a modular form of weight -6 for ρ_{D_K} . The components of F are given by

$$\begin{aligned} F_0 &= \frac{1}{\eta_g} + \frac{1}{2} \left(\frac{\theta_{0_0}}{\Delta} - \frac{1}{\eta_g} \right) \\ &= q^{-1} + 12 + 300q + 5792q^2 + 84186q^3 + 949920q^4 + 8813768q^5 + \dots \end{aligned}$$

and

$$\begin{aligned} F_\gamma &= \frac{1}{2} \left(\frac{\theta_{0_0}}{\Delta} - \frac{1}{\eta_g} \right) \\ &= 12 + 288q + 5792q^2 + 84096q^3 + 949920q^4 + 8813248q^5 + \dots \end{aligned}$$

if $\gamma \in D_K^2 \setminus \{0\}$,

$$F_\gamma = \frac{\theta_{0_L}}{2\Delta} = 4 + 224q + 5344q^2 + 81792q^3 + 939232q^4 + 8769856q^5 + \dots$$

if $\gamma^2/2 = 0 \pmod{1}$ and $\gamma \notin D_K^2$,

$$F_\gamma = \frac{\theta_{2_S}}{2\Delta} = q^{-1/2} + 44q^{1/2} + 1242q^{3/2} + 22216q^{5/2} + 287463q^{7/2} + \dots$$

if $\gamma^2/2 = 1/2 \pmod{1}$ and $\gamma + H \in 2_{L_S}$,

$$F_\gamma = \frac{\theta_{2_L}}{2\Delta} = 32q^{1/2} + 1152q^{3/2} + 21696q^{5/2} + 284928q^{7/2} + \dots$$

if $\gamma^2/2 = 1/2 \pmod{1}$ and $\gamma + H \in 2_S \cup 2_{L_L}$,

$$F_\gamma = \frac{\theta_3}{2\Delta} = q^{-1/4} + 90q^{3/4} + 2535q^{7/4} + 42614q^{11/4} + 521235q^{15/4} + \dots$$

if $\gamma^2/2 = 1/4 \pmod{1}$ and

$$F_\gamma = \frac{\theta_1}{2\Delta} = 12q^{1/4} + 520q^{5/4} + 10908q^{9/4} + 153960q^{13/4} + 1669720q^{17/4} + \dots$$

if $\gamma^2/2 = 3/4 \pmod{1}$.

The Fourier coefficients of the F_γ are nonnegative rational integers.

Now let

$$L = K \oplus II_{1,1}.$$

We can consider F as a modular form for ρ_{D_L} . Then the poles of F define a reflection group W of L' which has a Weyl vector. The roots of W are the vectors

$$\begin{aligned} \alpha &\in L \text{ with } \alpha^2 = 2, \\ \alpha &\in L' \text{ with } \alpha^2 = 1 \text{ and } (\alpha_K + K) + H \in 2_{L_S}, \\ \alpha &\in L' \text{ with } \alpha^2 = 1/2. \end{aligned}$$

The reflection group W has a Weyl vector ρ in $\mathbb{R} \otimes L$, i.e. the simple roots of W are the roots α satisfying $\rho\alpha = -\alpha^2/2$. The vector 2ρ is a primitive norm 0 vector in L' and $2\rho + L$ is in $D_L^2 \setminus \{0\}$.

Let

$$M = L \oplus II_{1,1}.$$

Then F defines a modular form for ρ_{D_M} . Let Ψ be the theta lift of F . The lattice M has one orbit of primitive norm 0 vectors of level 1 under $O(M)$ and has no primitive norm 0 vectors of level 4.

Theorem 6.4

The automorphic form Ψ is holomorphic and has singular weight 6. The level 1 expansion is given by

$$e((\rho, Z)) \prod_{\alpha \in L'^+} (1 - e((\alpha, Z)))^{[F_{\alpha+L}](-\alpha^2/2)} = \sum_{w \in W} \det(w) \eta_g((w\rho, Z))$$

As in the previous case this identity is also the denominator identity of a generalized Kac-Moody algebra whose real simple roots are the simple roots of W and imaginary simple roots are the positive integral multiples of 2ρ with multiplicity 12. The root lattice is L' and the multiplicity of a root α in L' is given by $[F_{\alpha+L}](-\alpha^2/2)$. This Lie algebra can also be constructed by orbifolding the fake monster algebra with a lift of g .

We show now that the Niemeier lattice N with root system D_{12}^2 gives a simpler construction of the above objects. The quotient D'_{12}/D_{12} can be represented by the elements $0 = (0^{12})$, $s = ((\frac{1}{2})^{12})$, $c = ((\frac{1}{2})^{11}(-\frac{1}{2}))$ and $v = (0^{11}1)$. Adding the glue vectors $(0, 0)$, (s, s) , (c, v) and (v, c) to the orthogonal sum $D_{12} \oplus D_{12}$ we obtain an even unimodular lattice N with root system D_{12}^2 . The direct product $O(D_{12}^+)^2$ is a normal subgroup of $O(N)$ with quotient S_2 . Define $g \in O(N)$ by $g(x, y) = (y, x)$. Then g has order 2 and N^g and $N^{g\perp}$ are both isomorphic to $\sqrt{2}D_{12}^+$. The centralizer $C(g)$ of g in $O(N)$ induces the full orthogonal group of N^g . Hence the orbits of the corresponding size are mapped into each other under the natural map $i : D_{N^g} \rightarrow D_{N^{g\perp}}$. We construct a modular form F as above. Then the components of F depend only on the orbits of $O(\sqrt{2}D_{12})$ on $D_{\sqrt{2}D_{12}}$. In particular the generators of the Weyl group of norm 1 correspond to the elements in the two orbits of length 132 in D_K .

Cycle shape 3.21

Let g be an automorphism of the Leech lattice of cycle shape 3.21. Then the fixed point lattice Λ^g has genus $II_{2,0}(3^{+27}7^{-1})$ and a Gram matrix is given by $(\begin{smallmatrix} 6 & 3 \\ 3 & 12 \end{smallmatrix})$. The lattice Λ^g represents the unique class in this genus. The centralizer of g induces the full orthogonal group of Λ^g (cf. also [HL]).

The lattice $\Lambda^{g,3} = \Lambda^{g^3} \cap \Lambda^{g\perp}$ has some very nice properties. It is isomorphic

to $\begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix} \otimes A_2$ and a Gram matrix is given by

$$\begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix} \otimes \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 4 & 2 & 2 & 1 \\ 2 & 4 & 1 & 2 \\ 2 & 1 & 8 & 4 \\ 1 & 2 & 4 & 8 \end{pmatrix}.$$

The lattice $\Lambda^{g,3}$ has genus $II_{4,0}(3^{+2}7^{-2})$. It is strongly modular, i.e. invariant up to isomorphism under the Atkin-Lehner involutions W_d , $d|21$. There are 3 classes in the genus of $\Lambda^{g,3}$ with minima 2, 2 and 4, i.e. $\Lambda^{g,3}$ represents the unique class without roots. The orthogonal group of $\Lambda^{g,3}$ has 40 orbits on the discriminant form $D_{\Lambda^{g,3}}$ and 4 orbits on $D_{\Lambda^{g,3},3}$. They are described in the following table.

| norm | length | order | name |
|------|--------|-------|-------|
| 0 | 1 | 1 | 0_0 |
| 1/3 | 4 | 3 | 1 |
| 2/3 | 2 | 3 | 2_S |
| | 2 | 3 | 2_L |

The corresponding theta functions are given by

$$\begin{aligned} \theta_{0_0}(\tau) &= 1 + 6q^2 + 12q^4 + 12q^5 + 6q^6 + 24q^7 + 18q^8 + 24q^{10} + \dots \\ \theta_1(\tau) &= 3q^{4/3} + 6q^{7/3} + 6q^{10/3} + 6q^{13/3} + 12q^{16/3} + 6q^{19/3} + \dots \\ \theta_{2_S}(\tau) &= 3q^{2/3} + 3q^{8/3} + 6q^{11/3} + 18q^{14/3} + 12q^{17/3} + 12q^{20/3} + \dots \\ \theta_{2_L}(\tau) &= 6q^{5/3} + 6q^{8/3} + 15q^{14/3} + 6q^{17/3} + 18q^{20/3} + 12q^{23/3} + \dots \end{aligned}$$

We have

$$(\theta_{2_S} - \theta_{2_L})/3\eta_{g^3}(\tau) = 1/\eta_g(\tau/3).$$

Since $\Lambda^{g,3}$ is a primitive sublattice of the Leech lattice the theta function of $\Lambda^{g,3}$ defines a modular form F for the Weil representation of the orthogonal complement $\Lambda^{g,3\perp}$ of $\Lambda^{g,3}$ in Λ . This lattice has genus $II_{20,0}(3^{+2}7^{-2})$. In particular there are no nontrivial isotropic elements in $D_{\Lambda^{g,3\perp},3}$. This implies that the function F cannot be written as a linear combination of liftings from $\Gamma_0(21)$ because such a function would be invariant under the automorphisms of $D_{\Lambda^{g,3\perp},3}$ but as we have seen above F does not have this symmetry.

We remark that $\Lambda^{g,7} = \Lambda^{g^7} \cap \Lambda^{g^\perp}$ is isomorphic to $\sqrt{3}A_6$ and has genus $II_{6,0}(3^{+6}7^{+1})$. There is exactly one class in this genus.

We write $\theta_{\Lambda^{g,3}}$ for the scalar valued theta function of $\Lambda^{g,3}$. Then $\theta_{\Lambda^{g,3}}$ is a modular form for $\Gamma_0(21)$ of weight 2 with trivial character. We describe the expansions at the cusps. Let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ such that $c|21$, $c > 0$ and $d = 0 \pmod{c'}$ where $c' = N/c$. Then

$$\theta_{\Lambda^{g,3}}|_M(\tau) = \epsilon \frac{1}{c'} \theta_{\Lambda^{g,3}}(\tau/c')$$

with $\epsilon = (\frac{c'}{2})$.

Let $K = \Lambda^g \oplus \sqrt{7}H_{1,1}$. Then K has genus $H_{3,1}(3^{+2}7^{+3})$. The quotient $\theta_{\Lambda^{g,3}}/\eta_{g^3}$ where $\eta_{g^3}(\tau) = \eta(\tau)^3\eta(7\tau)^3$ is a modular form for $\Gamma_0(21)$ of weight -1 with character $\chi(M) = (\frac{d}{7})$ for $M = (\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}) \in \Gamma_0(21)$. We have

$$\begin{aligned}\theta_{\Lambda^{g,3}}(\tau)/\eta_{g^3}(\tau) &= q^{-1} + 3 + 15q + 40q^2 + 117q^3 + 288q^4 + 677q^5 + \dots \\ \theta_{\Lambda^{g,3}}(\tau/3)/\eta_{g^3}(\tau) &= q^{-1} + 6q^{-1/3} + 3 + 12q^{1/3} + 30q^{2/3} + 15q + \dots\end{aligned}$$

The lift of $\theta_{\Lambda^{g,3}}/\eta_{g^3}$ on K with trivial support is given by

$$\begin{aligned}F_{\Gamma_0(21), \theta_{\Lambda^{g,3}}/\eta_{g^3}, 0} = \\ \frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} e^0 + \frac{1}{3} \sum_{\gamma \in D_{K,3}} g_{3,j_\gamma} e^\gamma + \sum_{\gamma \in D_{K,7}} g_{7,j_\gamma} e^\gamma + \frac{1}{3} \sum_{\gamma \in D_K} g_{21,j_\gamma} e^\gamma\end{aligned}$$

where the $g_{c,j}$ are T -eigenfunctions with eigenvalue $e(j/c)$ defined by the following decompositions

$$\begin{aligned}\theta_{\Lambda^{g,3}}(\tau/3)/\eta_{g^3}(\tau) &= g_{3,0}(\tau) + g_{3,1}(\tau) + g_{3,2}(\tau) \\ \theta_{\Lambda^{g,3}}(\tau/7)/\eta_{g^3}(\tau/7) &= g_{7,0}(\tau) + g_{7,1}(\tau) + \dots + g_{7,6}(\tau) \\ \theta_{\Lambda^{g,3}}(\tau/21)/\eta_{g^3}(\tau/7) &= g_{21,0}(\tau) + g_{21,1}(\tau) + \dots + g_{21,20}(\tau).\end{aligned}$$

Note that

$$\begin{aligned}\theta_{\Lambda^{g,3}}/\eta_{g^3} &= g_{3,0} \\ g_{7,j} &= g_{21,3j}.\end{aligned}$$

The functions with poles are $\theta_{\Lambda^{g,3}}/\eta_{g^3} = g_{3,0}$ and

$$\begin{aligned}g_{3,2} &= 6q^{-1/3} + 30q^{2/3} + 108q^{5/3} + 306q^{8/3} + 834q^{11/3} + \dots \\ g_{21,20} &= 6q^{-1/21} + 9792q^{20/21} + 835008q^{41/21} + 28697184q^{62/21} + \dots \\ g_{7,6} &= g_{21,18} = q^{-1/7} + 1494q^{6/7} + 143829q^{13/7} + 5254648q^{20/7} + \dots\end{aligned}$$

For $c|63$ we define the functions $h_c(\tau) = 1/\eta_g(\tau/c)$ and their T -eigenfunctions $h_{c,j}$ with $h_{c,j}|T = e(j/c)h_{c,j}$.

The group $D_{K,3}$ is generated by two orthogonal elements γ_1, γ_2 of norm $\gamma_1^2/2 = \gamma_2^2/2 = 1/3 \pmod{1}$. The function h_3 is a modular form of weight -1 for $\Gamma_1(21)$ with character $\chi(M) = e(-b/3)$ for $M = (\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}) \in \Gamma_1(21)$. The lift of h_3 on γ_1 is given by (cf. Theorem 3.7)

$$\begin{aligned}\frac{1}{12}F_{\Gamma_1(21), h_3, \gamma_1} &= h_3(e^{\gamma_1} + e^{-\gamma_1}) - h_3(e^{\gamma_2} + e^{-\gamma_2}) \\ &\quad - \sum_{\gamma \in \gamma_1 + D_{K,7}} h_{21,j_\gamma}(e^\gamma + e^{-\gamma}) + \sum_{\gamma \in \gamma_2 + D_{K,7}} h_{21,j_\gamma}(e^\gamma + e^{-\gamma}).\end{aligned}$$

We have

$$\begin{aligned}
& \frac{1}{4}F_{\Gamma_0(21),\theta_{\Lambda^g,3}/\eta_{g^3},0} + \frac{1}{24}F_{\Gamma_1(21),h_3,\gamma_1} \\
&= \frac{1}{4} \frac{\theta_{\Lambda^g,3}}{\eta_{g^3}} e^0 \\
&+ \frac{1}{12} \sum_{\substack{\gamma \in D_{K,3} \\ \gamma^2/2 \not\equiv 1/3 \pmod{1}}} g_{3,j_\gamma} e^\gamma \\
&+ \frac{1}{12} (g_{3,2} + 6h_3)(e^{\gamma_1} + e^{-\gamma_1}) \\
&+ \frac{1}{12} (g_{3,2} - 6h_3)(e^{\gamma_2} + e^{-\gamma_2}) \\
&+ \frac{1}{4} \sum_{\gamma \in D_{K,7}} g_{7,j_\gamma} e^\gamma \\
&+ \frac{1}{12} \sum_{\substack{\gamma \in D_K \\ \gamma^2/2 \not\equiv 1/3+n/7 \pmod{1}}} g_{21,j_\gamma} e^\gamma \\
&+ \frac{1}{12} \sum_{\gamma \in \gamma_1 + D_{K,7}} (g_{21,j_\gamma} - 6h_{21,j_\gamma})(e^\gamma + e^{-\gamma}) \\
&+ \frac{1}{12} \sum_{\gamma \in \gamma_2 + D_{K,7}} (g_{21,j_\gamma} + 6h_{21,j_\gamma})(e^\gamma + e^{-\gamma}).
\end{aligned}$$

The Fourier expansions of the functions $g_{3,1}/12$, $(g_{3,2} \pm 6h_3)/12$ and $g_{21,3j}/12$, $g_{21,3j+1}/12$, $(g_{21,3j+2} \pm 6h_{21,3j+2})/12$ have nonnegative integral coefficients, for example

$$\begin{aligned}
\frac{1}{12} g_{3,1} &= q^{1/3} + 5q^{4/3} + 17q^{7/3} + 48q^{10/3} + 123q^{13/3} + 286q^{16/3} + \dots \\
&= \frac{\theta_1}{3\eta_{g^3}} \\
\frac{1}{12} (g_{3,2} + 6h_3) &= q^{-1/3} + 3q^{2/3} + 10q^{5/3} + 27q^{8/3} + 72q^{11/3} + 170q^{14/3} + \dots \\
&= \frac{\theta_{2_S}}{3\eta_{g^3}} \\
\frac{1}{12} (g_{3,2} - 6h_3) &= 2q^{2/3} + 8q^{5/3} + 24q^{8/3} + 67q^{11/3} + 163q^{14/3} + 375q^{17/3} + \dots \\
&= \frac{\theta_{2_L}}{3\eta_{g^3}}.
\end{aligned}$$

Let $M = \Lambda^g \oplus \sqrt{7}H_{1,1} \oplus H$ where H is a lattice with Gram matrix $\begin{pmatrix} -2 & 3 \\ 3 & 0 \end{pmatrix}$. Then M has genus $II_{4,2}(3^{+2}9^{-1}7^{+3})$.

The modular form

$$\frac{1}{4} F_{\Gamma_0(21), \theta_{\Lambda^g, 3}/\eta_{g^3}, 0} + \frac{1}{24} F_{\Gamma_1(21), h_3, \gamma_1}$$

on K induces a modular form $F_{\theta_{\Lambda^g, 3}/\eta_{g^3}}$ on M .

We remark that $F_{\theta_{\Lambda^g, 3}/\eta_{g^3}}$ can also be obtained as

$$F_{\theta_{\Lambda^g, 3}/\eta_{g^3}} = \frac{1}{12} F_{\Gamma_0(63), \theta_{\Lambda^g, 3}/\eta_{g^3}, D^{21}} + \frac{1}{216} \sum_{\gamma \in \gamma_1 + D^{21}} F_{\Gamma_1(63), h_3, \gamma}.$$

The function $\eta_g(\tau) = \eta(3\tau)\eta(21\tau)$ is a modular form for $\Gamma_0(63)$ of weight 1 with character $\chi(M) = \left(\frac{d}{7}\right)$ for $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(63)$. For the lift of $1/\eta_g$ on M with trivial support we find

$$\begin{aligned} & \frac{1}{3} F_{\Gamma_0(63), 1/\eta_g, 0} \\ &= \frac{1}{\eta_g} e^0 - \frac{1}{3} \sum_{\gamma \in D^{21}} \frac{1}{\eta_g} e^\gamma + \sum_{\gamma \in D_7} h_{7, j_\gamma} e^\gamma - \frac{1}{3} \sum_{\gamma \in D^3} h_{7, j_\gamma} e^\gamma \\ & \quad + \sum_{\gamma \in D_9} h_{9, j_\gamma} e^\gamma + \sum_{\gamma \in D} h_{63, j_\gamma} e^\gamma. \end{aligned}$$

Let

$$F = F_{\theta_{\Lambda^g, 3}/\eta_{g^3}} + \frac{1}{3} F_{\Gamma_0(63), 1/\eta_g, 0}.$$

Then F is a modular form of weight -1 on M . Using $\theta_{\Lambda^g, 3}/\eta_{g^3} = g_{3,0}$ and $g_{7,j} = g_{21,3j}$ we obtain

Theorem 6.5

The function F is given by

$$\begin{aligned}
F &= \frac{1}{\eta_g} e^0 + \frac{1}{3} \sum_{\gamma \in D^{21}} \left(\frac{\theta_{\Lambda_{g,3}}}{\eta_{g^3}} - \frac{1}{\eta_g} \right) e^\gamma \\
&+ \frac{1}{12} \sum_{\substack{\gamma \in D_3 \\ \gamma^2/2 = 2/3 \pmod{1}}} g_{3,1} e^\gamma \\
&+ \frac{1}{12} \sum_{\gamma \in \gamma_1 + D^{21}} (g_{3,2} + 6h_3) (e^\gamma + e^{-\gamma}) \\
&+ \frac{1}{12} \sum_{\gamma \in \gamma_2 + D^{21}} (g_{3,2} - 6h_3) (e^\gamma + e^{-\gamma}) \\
&+ \sum_{\gamma \in D_7} h_{7,j_\gamma} e^\gamma + \frac{1}{3} \sum_{\gamma \in D^3} (g_{7,j_\gamma} - h_{7,j_\gamma}) e^\gamma \\
&+ \frac{1}{12} \sum_{\substack{\gamma \in D_{21} \\ \gamma^2/2 = 2/3+n/7 \pmod{1}}} g_{21,j_\gamma} e^\gamma \\
&+ \frac{1}{12} \sum_{\gamma \in \gamma_1 + D^3} (g_{21,j_\gamma} - 6h_{21,j_\gamma}) (e^\gamma + e^{-\gamma}) \\
&+ \frac{1}{12} \sum_{\gamma \in \gamma_2 + D^3} (g_{21,j_\gamma} + 6h_{21,j_\gamma}) (e^\gamma + e^{-\gamma}) \\
&+ \sum_{\gamma \in D_9} h_{9,j_\gamma} e^\gamma \\
&+ \sum_{\gamma \in D} h_{63,j_\gamma} e^\gamma
\end{aligned}$$

The coefficients of the components F_γ of F are nonnegative rational integers.

Note that without the contributions of the lifting of h_3 the Fourier coefficients of the principal part of F would be rational but not integral so that we could not apply the singular theta correspondence to F .

We describe some components of F . They are important to understand the

properties of the Weyl group and the Weyl vector we define below.

Let γ be an isotropic element in D . Then γ is in $D^3 = D^{21} \oplus D_7$. We have

$$\begin{aligned}
F_\gamma &= \frac{1}{\eta_g} + \frac{1}{3} \left(\frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} - \frac{1}{\eta_g} \right) + h_{7,0} + \frac{1}{3} (g_{7,0} - h_{7,0}) \\
&= \frac{1}{\eta_g} + \frac{1}{3} \left(\frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} - \frac{1}{\eta_g} \right) + \frac{1}{7} \left(\frac{\theta_{\Lambda^{g,7}}}{\eta_{g^7}} - \frac{1}{\eta_g} \right) \\
&\quad + \frac{1}{21} \left(\frac{\theta_{\Lambda^{g^\perp}}}{\Delta} - \frac{\theta_{\Lambda^{g,7}}}{\eta_{g^7}} - \frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} + \frac{1}{\eta_g} \right) \\
&= q^{-1} + 2 + 1060q + 83728q^2 + 2790636q^3 + 57148320q^4 + \dots
\end{aligned}$$

if $\gamma = 0$,

$$\begin{aligned}
F_\gamma &= \frac{1}{3} \left(\frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} - \frac{1}{\eta_g} \right) + \frac{1}{3} (g_{7,0} - h_{7,0}) \\
&= \frac{1}{3} \left(\frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} - \frac{1}{\eta_g} \right) + \frac{1}{21} \left(\frac{\theta_{\Lambda^{g^\perp}}}{\Delta} - \frac{\theta_{\Lambda^{g,7}}}{\eta_{g^7}} - \frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} + \frac{1}{\eta_g} \right) \\
&= 2 + 1060q + 83720q^2 + 2790636q^3 + 57148320q^4 + 846724518q^5 + \dots
\end{aligned}$$

if $\gamma \in D^{21} \setminus \{0\}$,

$$\begin{aligned}
F_\gamma &= h_{7,0} + \frac{1}{3} (g_{7,0} - h_{7,0}) \\
&= \frac{1}{7} \left(\frac{\theta_{\Lambda^{g,7}}}{\eta_{g^7}} - \frac{1}{\eta_g} \right) + \frac{1}{21} \left(\frac{\theta_{\Lambda^{g^\perp}}}{\Delta} - \frac{\theta_{\Lambda^{g,7}}}{\eta_{g^7}} - \frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} + \frac{1}{\eta_g} \right) \\
&= 1 + 1055q + 83714q^2 + 2790597q^3 + 57148224q^4 + 846724377q^5 + \dots
\end{aligned}$$

if $\gamma \in D_7 \setminus \{0\}$, and

$$\begin{aligned}
F_\gamma &= \frac{1}{3} (g_{7,0} - h_{7,0}) \\
&= \frac{1}{21} \left(\frac{\theta_{\Lambda^{g^\perp}}}{\Delta} - \frac{\theta_{\Lambda^{g,7}}}{\eta_{g^7}} - \frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} + \frac{1}{\eta_g} \right) \\
&= 1 + 1055q + 83707q^2 + 2790597q^3 + 57148224q^4 + 846724293q^5 + \dots
\end{aligned}$$

if $\gamma \in D^3 \setminus \{D^{21} \cup D_7\}$.

Next we describe the singular components of F . They are given by

$$\begin{aligned}
F_0 &= \frac{1}{\eta_g} + \frac{1}{3} \left(\frac{\theta_{\Lambda^{g,3}}}{\eta_{g^3}} - \frac{1}{\eta_g} \right) + h_{7,0} + \frac{1}{3} (g_{7,0} - h_{7,0}) \\
&= q^{-1} + 2 + 1060q + 83728q^2 + 2790636q^3 + 57148320q^4 + \dots
\end{aligned}$$

and

$$\begin{aligned} F_\gamma &= \frac{1}{12}(g_{3,2} + 6h_3) + \frac{1}{12}(g_{21,14} - 6h_{21,14}) \\ &= q^{-1/3} + 166q^{2/3} + 22071q^{5/3} + 925654q^{8/3} + 21777668q^{11/3} + \dots \end{aligned}$$

if $\gamma^2/2 = 1/3 \pmod{1}$ and $\gamma \in \pm\gamma_1 + D^{21}$,

$$\begin{aligned} F_\gamma &= h_{7,6} + \frac{1}{3}(g_{7,6} - h_{7,6}) + h_{63,54} \\ &= h_{7,6} + \frac{1}{3}(g_{7,6} - h_{7,6}) \\ &= q^{-1/7} + 498q^{6/7} + 47943q^{13/7} + 1751560q^{20/7} + 37970952q^{27/7} + \dots \end{aligned}$$

if $\gamma^2/2 = 1/7 \pmod{1}$ and $\gamma \in D_7$,

$$\begin{aligned} F_\gamma &= h_{9,8} + h_{63,56} \\ &= q^{-1/9} + 584q^{8/9} + 54268q^{17/9} + 1943680q^{26/9} + 41603422q^{35/9} + \dots \end{aligned}$$

if $\gamma^2/2 = 1/9 \pmod{1}$ and $\gamma \in D_9$,

$$\begin{aligned} F_\gamma &= \frac{1}{12}(g_{21,20} + 6h_{21,20}) + h_{63,60} \\ &= \frac{1}{12}(g_{21,20} + 6h_{21,20}) \\ &= q^{-1/21} + 824q^{20/21} + 69660q^{41/21} + 2391912q^{62/21} + \dots \end{aligned}$$

if $\gamma^2/2 = 1/21 \pmod{1}$ and $\gamma \in \pm\gamma_2 + D^3$, and

$$\begin{aligned} F_\gamma &= h_{63,62} \\ &= q^{-1/63} + 960q^{62/63} + 78660q^{125/63} + 2650432q^{188/63} + \dots \end{aligned}$$

if $\gamma^2/2 = 1/63 \pmod{1}$.

The lattices $\sqrt{7}II_{1,1} \oplus H$ and $II_{1,1} \oplus \sqrt{7}H$ are in the same genus and therefore isomorphic. This implies that we can consider F as a modular form on

$$L = \Lambda^g \oplus \sqrt{7}H.$$

The poles of F define a reflection group W of L' . The roots of W are the vectors

$$\begin{aligned} &\alpha \in L \text{ with } \alpha^2 = 2, \\ &\alpha \in L' \text{ with } \alpha^2 = 2/3 \text{ and } \alpha + L \in \pm\gamma_1 + D_L^3, \\ &\alpha \in L' \text{ with } \alpha^2 = 2/7 \text{ and } \alpha + L \in D_{L,7}, \\ &\alpha \in L' \text{ with } \alpha^2 = 2/9 \text{ and } \alpha + L \in D_{L,9}, \\ &\alpha \in L' \text{ with } \alpha^2 = 2/21 \text{ and } \alpha + L \in \pm\gamma_2 + D_L^3, \\ &\alpha \in L' \text{ with } \alpha^2 = 2/63. \end{aligned}$$

The reflection group W has a Weyl vector ρ in $\mathbb{R} \otimes L$, i.e. the simple roots of W are the roots α satisfying $\rho\alpha = -\alpha^2/2$. The vector 3ρ is a primitive norm 0 vector in L' and $3\rho + L$ is in $D_L^3 \setminus (D_L^{21} \cup D_{L,7})$. There is one orbit of primitive norm 0 vectors in $(\sqrt{7}H)'$ under the orthogonal group of $(\sqrt{7}H)'$ and we can take any vector in this orbit for 3ρ .

The function F is a modular form for $M = L \oplus II_{1,1}$. Let Ψ be the theta lift of F . The lattice M has one orbit of primitive norm 0 vectors of level 1 under $O(M)$.

Theorem 6.6

The automorphic form Ψ is holomorphic and has singular weight 1. The level 1 expansion is given by

$$e((\rho, Z)) \prod_{\alpha \in L'^+} (1 - e((\alpha, Z)))^{[F_{\alpha+L}](-\alpha^2/2)} = \sum_{w \in W} \det(w) \eta_g((w\rho, Z))$$

As in the previous cases this identity is also the denominator identity of a generalized Kac-Moody algebra whose real simple roots are the simple roots of W and imaginary simple roots are the positive integral multiples $3n\rho$ of 3ρ with multiplicity 2 if $7|n$ and multiplicity 1 otherwise. The root lattice is L' and the multiplicity of a root α in L' is given by $[F_{\alpha+L}](-\alpha^2/2)$. Again this Lie algebra can also be constructed by orbifolding the fake monster algebra with a lift of g .

References

- [Ba] A. G. Barnard, *The singular theta correspondence, Lorentzian lattices and Borcherds-Kac-Moody algebras*, Ph.D. thesis, UC Berkeley, arXiv: math/0307102 (2003)
- [B1] R. E. Borcherds, *Monstrous moonshine and monstrous Lie superalgebras*, Invent. Math. **109** (1992), 405–444
- [B2] R. E. Borcherds, *Automorphic forms with singularities on Grassmannians*, Invent. Math. **132** (1998), 491–562
- [BCP] W. Bosma, J. Cannon and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), 235–265
- [Br] J. H. Bruinier, *Borcherds products on $O(2, l)$ and Chern classes of Heegner divisors*, Lecture Notes in Mathematics **1780**, Springer, Berlin, 2002
- [CS] J. H. Conway, N. J. A. Sloane, *Sphere packings, lattices and groups*, 3rd ed., Grundlehren der mathematischen Wissenschaften **290**, Springer, New York, 1999
- [DS] F. Diamond, J. Shurman, *A first course in modular forms*, Graduate Texts in Mathematics, Springer, New York, 2005

- [HL] K. Harada, M.-L. Lang, *On some sublattices of the Leech lattice*, Hokkaido Math. J. **19** (1990), 435–446
- [N] V. V. Nikulin, *Integer symmetric bilinear forms and some of their geometric applications*, Math. USSR Izv. **14** (1979), 103–167
- [O] T. O. O’Meara, *Introduction to quadratic forms*, Classics in Mathematics, Springer, Berlin, 2000
- [P] The PARI Group, Bordeaux, PARI/GP, version 2.3.4, 2008, available from <http://pari.math.u-bordeaux.fr/>
- [S1] N. R. Scheithauer, *On the classification of automorphic products and generalized Kac-Moody algebras*, Invent. Math. **164** (2006), 641–678
- [S2] N. R. Scheithauer, *The Weil representation of $SL_2(\mathbb{Z})$ and some applications*, Int. Math. Res. Notices 2009, no. 8, 1488–1545