

# Bounded Imaginary Powers and $H_\infty$ -Calculus of the Stokes Operator in Two-Dimensional Exterior Domains

Helmut Abels\*

September 1, 2004

## Abstract

The present contribution deals with the Stokes operator  $A_q$  on  $L_\sigma^q(\Omega)$ ,  $1 < q < \infty$ , where  $\Omega$  is an exterior domain in  $\mathbb{R}^2$  of class  $C^2$ . It is proved that  $A_q$  admits a bounded  $H_\infty$ -calculus. This implies the existence of bounded imaginary powers of  $A_q$ , which has several important applications. – So far this property was only known for exterior domains in  $\mathbb{R}^n$ ,  $n \geq 3$ . – In particular, this shows that  $A_q$  has maximal regularity on  $L_\sigma^q(\Omega)$ . For the proof the resolvent  $(\lambda + A_q)^{-1}$  has to be analyzed for  $|\lambda| \rightarrow \infty$  and  $\lambda \rightarrow 0$ . For large  $\lambda$  this is done using an approximate resolvent based on the results of [3], which were obtained by applying the calculus of pseudodifferential boundary value problems. For small  $\lambda$  we analyze the representation of the resolvent developed in [11] by a potential theoretical method.

**Key words:** Stokes operator, Stokes equations, exterior domains, bounded imaginary powers,  $H_\infty$ -calculus

**AMS-Classification:** 35 Q 30, 76 D 07, 47 A 60, 47 F 05

## 1 Introduction and Main Result

Let  $\Omega \subset \mathbb{R}^2$  be an exterior domain with  $C^2$ -boundary, i.e.,  $\mathbb{R}^2 \setminus \Omega$  is compact and  $\partial\Omega$  is a  $C^2$ -manifold. Moreover, let  $L_\sigma^q(\Omega) := \overline{\{f \in C_0^\infty(\Omega)^n : \operatorname{div} f = 0\}}^{L^q(\Omega)}$ ,  $1 < q < \infty$ , denote the space of solenoidal vector fields in  $L^q(\Omega)^n$  with vanishing normal component on  $\partial\Omega$ .

In this article we consider the Stokes operator  $A_q = -P_q \Delta$  on  $L_\sigma^q(\Omega)$  with domain

$$\mathcal{D}(A_q) = \{f \in W_q^2(\Omega)^n : f|_{\partial\Omega} = 0\} \cap L_\sigma^q(\Omega)$$

---

\*Department of Mathematics, Darmstadt University of Technology, Schloßgartenstr. 7, 64289 Darmstadt, Germany, e-mail: abels@mathematik.tu-darmstadt.de

where  $P_q : L^q(\Omega)^n \rightarrow L^q_\sigma(\Omega)$  denotes the well-known Helmholtz projection, cf. Simader and Sohr [23]. Borchers and Varnhorn [11] proved that  $-A_q$  generates a bounded and analytic semi-group. More precisely, they have shown that

$$\|(\lambda + A_q)^{-1}\|_{\mathcal{L}(L^q_\sigma(\Omega))} \leq \frac{C_{q,\delta}}{|\lambda|}, \quad \lambda \in \Sigma_\delta, \quad (1.1)$$

where  $\Sigma_\delta := \{z \in \mathbb{C} \setminus \{0\} : |\arg z| < \delta\}$  and  $\delta \in (0, \pi)$ . Earlier the same property was shown by Borchers and Sohr [10] for the case of exterior domains  $\Omega \subset \mathbb{R}^n$  with  $n \geq 3$ . But the latter contribution could not settle the two-dimensional case since there are some additional difficulties in comparison to the case  $n \geq 3$ , cf. Remark 1.3 below and [11, Introduction] for further explanations.

Besides the fact that  $-A_q$  generates a bounded analytic semi-group, an important property of the Stokes operator is that it possesses *bounded imaginary powers*, i.e.,

$$A_q^{iy} := \frac{1}{2\pi i} \int_\Gamma (-\lambda)^{iy} (\lambda + A_q)^{-1} d\lambda$$

is a bounded operator satisfying

$$\|A_q^{iy}\|_{\mathcal{L}(L^q_\sigma(\Omega))} \leq C_{q,\delta} e^{(\pi-\delta)|y|}, \quad y \in \mathbb{R}, \quad (1.2)$$

where  $\delta \in (0, \pi)$ ,  $1 < q < \infty$ , and  $\Gamma$  is the negatively orientated boundary of  $\Sigma_\delta$ . The latter property is not difficult to prove in the case  $\Omega = \mathbb{R}^n$  or  $\Omega = \mathbb{R}^n_+$ ,  $n \geq 2$ , cf. [18]. Besides these cases, the proof (1.2) is involved and most proofs use pseudodifferential operator techniques. This was done by Giga [16] for bounded domains in  $\mathbb{R}^n$ ,  $n \geq 2$ , and by Giga and Sohr [17] for exterior domains in  $\mathbb{R}^n$ ,  $n \geq 3$ , with smooth boundary. An alternative proof, which uses a perturbation theorem for the  $H_\infty$ -calculus instead of pseudodifferential operators, was given by Noll and Saal [22] for bounded and exterior domains in  $\mathbb{R}^n$ ,  $n \geq 3$ , with  $C^3$ -boundary. Moreover, (1.2) was proved for an infinite layer  $\Omega = \mathbb{R}^{n-1} \times (-1, 1)$  in [4] and more generally for so-called asymptotically flat layers with  $C^{1,1}$ -boundary in [3].

The purpose of the present contribution is to prove:

**Theorem 1.1** *Let  $1 < q < \infty$  and let  $\delta \in (0, \pi)$ . Then  $A_q$  admits a bounded  $H_\infty$ -calculus with respect to  $\delta$ , i.e.,*

$$h(A_q) := \frac{1}{2\pi i} \int_\Gamma h(-\lambda) (\lambda + A_q)^{-1} d\lambda \quad (1.3)$$

*is a bounded operator satisfying*

$$\|h(A_q)\|_{\mathcal{L}(L^q_\sigma(\Omega))} \leq C_{q,\delta} \|h\|_\infty \quad (1.4)$$

*for all  $h \in H_\infty(\delta)$ , where  $H_\infty(\delta)$  denotes the Banach algebra of all bounded holomorphic functions  $h : \Sigma_{\pi-\delta} \rightarrow \mathbb{C}$ .*

We note that in order to prove (1.4) for all  $h \in H_\infty(\delta)$  it is sufficient to show the estimate for  $h \in H(\delta)$ , which consists of all  $h \in H_\infty(\delta)$  such that

$$|h(z)| \leq C \frac{|z|^s}{1 + |z|^{2s}} \quad \text{for all } z \in \Sigma_{\pi-\delta}$$

for some  $s > 0$ , cf. [7, Lemma 2.1]. For  $h \in H(\delta)$  the integral (1.3) is well-defined as a Bochner integral on  $\mathcal{L}(L_\sigma^q(\Omega))$  and for arbitrary  $h \in H_\infty(\delta)$  the operator in (1.3) can be defined on a suitable dense subspace, cf. [7] for details.

The bounded  $H_\infty$ -calculus was introduced by McIntosh [21] and generalizes the property of having bounded imaginary powers since choosing  $h_y(z) := z^{iy}$ ,  $y \in \mathbb{R}$ , in (1.3)-(1.4) implies (1.2). Although in [4, 16, 17, 18] only (1.2) is proved, the proofs are easily modified to show (1.4).

The well-known result due to Dore and Venni [14, Theorem 3.2] and its extension by Giga and Sohr [18, Theorem 2.1] gives an important application of this abstract property:

**Theorem 1.2** *Let  $1 < p, q < \infty$  and let  $0 < T \leq \infty$ . Then for every  $f \in L^p(0, T; L_\sigma^q(\Omega))$  there is a unique solution  $u \in W_p^1(0, T; L_\sigma^q(\Omega)) \cap L^p(0, T; \mathcal{D}(A_q))$  of*

$$\begin{aligned} u'(t) + A_q u(t) &= f(t), & 0 < t < T, \\ u(0) &= 0 \end{aligned}$$

Moreover,

$$\|u'\|_{L^p(0, T; L_\sigma^q)} + \|A_q u\|_{L^p(0, T; L_\sigma^q)} \leq C \|f\|_{L^p(0, T; L_\sigma^q)},$$

where  $C$  does not depend on  $T$ .

Therefore the Stokes operator  $A_q$  has *maximal regularity* on  $L_\sigma^q(\Omega)$ ,  $1 < q < \infty$ .

Finally, we mention that the boundedness of  $A^{iy}$  and (1.2) can be used to characterize the domain of the fractional powers  $A_q^\alpha$ ,  $0 < \alpha < 1$ , as

$$\mathcal{D}(A_q^\alpha) = (L_\sigma^q(\Omega), \mathcal{D}(A_q))_{[\alpha]},$$

where  $(\cdot, \cdot)_{[\alpha]}$  denotes the complex interpolation functor, cf. [17, Proposition 6.1].

The outline of the proof of Theorem 1.1 is as follows: Roughly speaking, one has to deal with two singularities of the Cauchy-integral (1.4). The first occurs since  $\Gamma$  is unbounded and  $(\lambda + A_q)^{-1}$  is only of order  $O(|\lambda|^{-1})$  as  $|\lambda| \rightarrow \infty$ . The second is due to the singularity of  $(\lambda + A_q)^{-1}$  as  $\lambda \rightarrow 0$ . Hence we split  $\Gamma = \Gamma_R \dot{\cup} \Gamma'_R$ , where  $\Gamma'_R := \Gamma \cap B_R(0)$  with suitable  $R > 0$  and analyze each part separately. In order to analyze (1.3) with  $\Gamma$  replaced by  $\Gamma_R$ , an approximate resolvent  $R_\lambda$  is constructed that coincides with  $(\lambda + A_q)^{-1}$  modulo terms of order  $O((1 + |\lambda|)^{-1-\varepsilon})$  for some  $\varepsilon > 0$ , as  $|\lambda| \rightarrow \infty$ . The construction is based on results of [3], which were obtained by pseudodifferential operator techniques and were used in order to prove (1.4) for

the Stokes operator on asymptotically flat layers. The latter analysis is done in Section 4. Finally, it remains to estimate (1.3) with  $\Gamma$  replaced by  $\Gamma'_R$ . For this we use the representation of  $(\lambda + A_q)^{-1}$  developed in [10] in terms of the resolvent of the Stokes operator on  $\mathbb{R}^2$  and some single and double layer potentials, cf. Section 5. But first of all we start with some preliminaries in Section 2 and introduce the so-called *reduced* Stokes operator in Section 3, which is needed in order to apply the results of [3].

**Remark 1.3** The analysis of the resolvent as  $|\lambda| \rightarrow 0$  in the present two-dimensional case is more difficult than in dimension  $n \geq 3$  since the estimate

$$\|\nabla^2 u\|_q \leq C_{q,\delta} \|A_q u\|_q, \quad \lambda \in \Sigma_\delta, u \in \mathcal{D}(A_q), \quad (1.5)$$

holds if and only if  $1 < q < \frac{n}{2}$ , cf. [11, Introduction] and [9]. Hence the method of [17, Section 4] for the case of an exterior domain in  $\mathbb{R}^n$ ,  $n \geq 3$ , is not applicable since it is based on (1.5).

**Acknowledgements:** The author is grateful to Hermann Sohr for pointing out that the present problem was unsolved before and for motivating him to solve it.

## 2 Preliminaries

First of all,  $\mathbb{N}$  will denote the set of natural numbers (without 0) and  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ .

Let  $\Omega \subseteq \mathbb{R}^n$ ,  $n \geq 2$ , be a domain. Then  $C^\infty(\Omega)$  denotes the set of all smooth  $f: \Omega \rightarrow \mathbb{C}$ ,  $C_0^\infty(\Omega)$  is the set of all  $f \in C^\infty(\Omega)$  with compact support, and

$$C_{(0)}^\infty(\overline{\Omega}) := \{u = v|_{\overline{\Omega}} : v \in C_0^\infty(\mathbb{R}^n)\}.$$

The usual Lebesgue-space with respect to the Lebesgue measure on  $\Omega$  and the  $(n-1)$ -dimensional surface measure on  $\partial\Omega$  will be denoted by  $L^q(\Omega)$ ,  $L^q(\partial\Omega)$ , resp.,  $1 \leq q \leq \infty$ . Moreover, we use the abbreviations  $\|\cdot\|_q \equiv \|\cdot\|_{L^q(\Omega)}$  and  $\|\cdot\|_{q,\partial\Omega} \equiv \|\cdot\|_{L^q(\partial\Omega)}$ . Furthermore,  $L_{\text{loc}}^q(\overline{\Omega})$ ,  $1 \leq q \leq \infty$ , is defined as the space of  $f: \Omega \rightarrow \mathbb{C}$  such that  $f \in L^q(B \cap \Omega)$  for all balls  $B$  with  $B \cap \Omega \neq \emptyset$ . The usual scalar product on  $L^2(M)$  is denoted by  $(\cdot, \cdot)_M$  for  $M = \Omega, \partial\Omega$ .

In the following the usual Sobolev-Slobodeckij spaces based on  $L^q$ ,  $1 < q < \infty$ , are denoted by  $W_q^s(\Omega)$  and  $W_q^s(\partial\Omega)$ ,  $s \geq 0$ , with norms  $\|\cdot\|_{s,q}$  and  $\|\cdot\|_{s,q,\partial\Omega}$ , respectively, cf. e.g. [6]. Moreover,  $W_{q,0}^m(\Omega)$ ,  $m \in \mathbb{N}$ , denotes the closure of  $C_0^\infty(\Omega)$  in  $W_q^m(\Omega)$  and

$$W_q^{-m}(\Omega) := (W_{q',0}^m(\Omega))', \quad W_{q,0}^{-m}(\Omega) := (W_{q'}^m(\Omega))', \quad W_q^{-s}(\partial\Omega) := (W_{q'}^s(\partial\Omega))'$$

for  $m \in \mathbb{N}$  and  $s > 0$ , where  $\frac{1}{q} + \frac{1}{q'} = 1$ .

Finally, the homogeneous Sobolev space of order 1 is defined as

$$\dot{W}_q^1(\Omega) := \{p \in L_{\text{loc}}^q(\overline{\Omega}) : \nabla p \in L^q(\Omega)\}$$

normed by  $\|\nabla \cdot\|_q$ . If  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , is an exterior domain with  $C^1$ -boundary, for every  $p \in \dot{W}_q^1(\Omega)$  there is a  $\tilde{p} \in \dot{W}_q^1(\mathbb{R}^n)$  such that  $\tilde{p}|_\Omega = p$  and  $\|\nabla \tilde{p}\|_q \leq C \|\nabla p\|_q$ , cf. e.g. [12, Theorem 1.2]. As a consequence we obtain the following useful lemma.

**Lemma 2.1** *Let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , be an exterior domain with  $C^1$ -boundary and let  $1 < q < \infty$ . Then for every  $p \in \dot{W}_q^1(\Omega)$  there is a decomposition  $p = p_1 + p_2$  such that  $p_1 \in W_q^1(\Omega)$  and  $p_2 \in L_{\text{loc}}^q(\overline{\Omega})$  with  $\nabla p_2 \in W_q^1(\Omega)$  satisfying  $\|(p_1, \nabla p_2)\|_{1,q} \leq C \|\nabla p\|_q$ .*

**Proof:** Define for instance  $p_1 \in W_q^1(\Omega)$  as

$$p_1 = \mathcal{F}^{-1}[(1 - \varphi)(\xi)\mathcal{F}[\tilde{p}](\xi)]|_{\Omega},$$

where  $\tilde{p}$  is as above,  $\varphi \in C_0^\infty(\mathbb{R}^n)$  with  $\varphi(\xi) = 1$  in a neighborhood of 0, and  $\mathcal{F}$  denotes the Fourier transformation.  $\blacksquare$

In the following let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , be an exterior domain with  $C^m$ -boundary,  $m \in \mathbb{N}$ . Concerning traces recall that, if  $m \geq s > j + \frac{1}{q}$ ,  $j \in \mathbb{N}_0$ , with  $s - \frac{1}{q} \notin \mathbb{N}$ , there is a bounded operator

$$\gamma_j : W_q^s(\Omega) \rightarrow W_q^{s-\frac{1}{q}}(\partial\Omega) \quad (2.1)$$

such that  $\gamma_j u = \partial_\nu^j u|_{\partial\Omega}$  for all  $f \in C_{(0)}^\infty(\overline{\Omega})$ , cf. e.g. [6].

Furthermore, we recall that for  $f \in L^q(\Omega)^n$  such that  $\text{div } f \in L^q(\Omega)$  it is possible to define a weak trace of the normal component  $\gamma_\nu f \in W_q^{-\frac{1}{q}}(\partial\Omega)$  by

$$\langle \gamma_\nu f, v \rangle_{\partial\Omega} := (f, \nabla v)_\Omega + (\text{div } f, v)_\Omega \quad \text{for all } v \in W_{q'}^1(\Omega). \quad (2.2)$$

Moreover, we note that, if  $f = f_0 + \nabla p$ ,  $f_0 \in L_\sigma^q(\Omega)$ ,  $p \in \dot{W}_q^1(\Omega)$ , is the Helmholtz decomposition of  $f \in L^q(\Omega)^n$ , then  $p$  is uniquely determined as solution of the weak Neumann problem

$$\Delta p = \text{div } f \quad \text{in } \Omega \quad (2.3)$$

$$\partial_\nu p|_{\partial\Omega} = \nu \cdot f|_{\partial\Omega} \quad \text{on } \partial\Omega, \quad (2.4)$$

where  $\nu$  denotes the exterior normal, (2.3) is understood in the sense of distributions, and (2.4) is understood as  $\gamma_\nu(f - \nabla p) = 0$ , cf. [23]. Because of the definition of  $\gamma_\nu$ , the pressure  $p \in \dot{W}_q^1(\Omega)$  solves (2.3)-(2.4) if and only if

$$(\nabla p, \nabla v)_\Omega = (f, \nabla v)_\Omega \quad \text{for all } v \in \dot{W}_{q'}^1(\Omega).$$

Finally, the resolvent of the Laplace operator on  $\mathbb{R}^n$ ,  $n \geq 2$ , is given by

$$(\lambda - \Delta_{\mathbb{R}^n})^{-1} f = \mathcal{F}^{-1} \left[ \frac{\mathcal{F}[f](\xi)}{\lambda + |\xi|^2} \right], \quad (2.5)$$

where  $\mathcal{F}$  denotes the Fourier transformation. As a consequence of the Mihlin multiplier theorem, cf. e.g. [8], one obtains for  $u = (\lambda - \Delta_{\mathbb{R}^n})^{-1} f$

$$|\lambda| \|u\|_q + \|\nabla^2 u\|_q \leq C_{q,\delta} \|f\|_q, \quad \lambda \in \Sigma_\delta, f \in L^q(\mathbb{R}^n), \quad (2.6)$$

where  $1 < q < \infty$  and  $0 < \delta < \pi$ .

### 3 The Reduced Stokes Operator

In the following let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , be an exterior domain with  $C^2$ -boundary.

In order to apply the results from [3] for the construction of an approximate resolvent for large  $\lambda$ , we need the *reduced* Stokes operator  $A_{0,q}$  defined as

$$A_{0,q}u := (-\Delta + \nabla K_N T)u, \quad Tu := \gamma_\nu(\Delta - \nabla \operatorname{div})u|_{\partial\Omega},$$

for  $u \in \mathcal{D}(A_{0,q}) := W_q^2(\Omega)^n \cap W_{q,0}^1(\Omega)^n$ , where  $K_N$  denotes the Poisson operator of the Laplace equation with Neumann boundary conditions, i.e.,  $\Delta K_N = 0$  and  $\partial_\nu K_N|_{\partial\Omega} = I$ . Because of [23, Theorem 4.4],  $K_N$  exists and is a bounded operator

$$K_N: W_{q,(0)}^{-\frac{1}{q}}(\partial\Omega) := \left\{ a \in W_q^{-\frac{1}{q}}(\partial\Omega) : \langle a, 1 \rangle_{\partial\Omega} = 0 \right\} \rightarrow \dot{W}_q^1(\Omega).$$

Note that  $A_{q,0}$  is a densely defined unbounded operator on  $L^q(\Omega)^n$  in contrast to the Stokes operator, which acts on the subspace  $L_q^q(\Omega)$ .

It remains to justify that  $\nabla K_N T: \mathcal{D}(A_{0,q}) \rightarrow L^q(\Omega)^n$ . Since  $\operatorname{div}(\Delta - \nabla \operatorname{div})u = 0$ ,  $T: W_q^2(\Omega)^n \rightarrow W_q^{-\frac{1}{q}}(\Omega)$  is a bounded operator and by (2.2)

$$\langle Tu, v \rangle_{\partial\Omega} := ((\Delta - \nabla \operatorname{div})u, \nabla v)_\Omega \quad \text{for all } v \in W_q^1(\Omega). \quad (3.1)$$

The latter identity implies that even  $Tu \in W_{q,(0)}^{-\frac{1}{q}}(\partial\Omega)$  for  $u \in W_q^2(\Omega)$ , which can be seen as follows: Since  $\partial\Omega$  is compact, we may assume that also  $\operatorname{supp} u$  is compact. Then choosing  $v \in C_{(0)}^\infty(\overline{\Omega})$  such that  $v \equiv 1$  on  $\operatorname{supp} u \cup \partial\Omega$  yields  $\langle Tu, 1 \rangle_{\partial\Omega} = 1$ .

There is the following alternative description of  $T$ : Introducing local coordinates  $Tu = \operatorname{div}_\tau \partial_\nu u_\tau|_{\partial\Omega}$  for every  $u \in C_{(0)}^\infty(\overline{\Omega})$  with  $u|_{\partial\Omega} = 0$ , cf. [20, Lemma A.1]. Here  $u_\tau$  denotes the tangential components of  $u$  and  $\operatorname{div}_\tau a_\tau := \partial_{\kappa_1} a_{\kappa_1} + \dots + \partial_{\kappa_{n-1}} a_{\kappa_{n-1}}$  for  $a \in W_q^s(\partial\Omega)$ ,  $s - \frac{1}{q} \notin \mathbb{Z}$ , where  $\kappa_1(y), \dots, \kappa_{n-1}(y)$  denotes a basis of the tangential space  $T_y \partial\Omega$  for each  $y \in \partial\Omega$  such that  $\kappa_1(y), \dots, \kappa_{n-1}(y), \nu(y)$  is a positively orientated orthonormal basis of  $\mathbb{R}^n$ . Hence by (2.1)

$$T: W_q^{2+s}(\Omega)^n \rightarrow W_q^{s-\frac{1}{q}}(\partial\Omega) \quad \text{for all } 0 \geq s > -1 + \frac{1}{q}. \quad (3.2)$$

**Remark 3.1** As the usual Stokes operator is associated to the Stokes resolvent equation, i.e.,

$$(\lambda - \Delta)u + \nabla p = f \quad \text{in } \Omega, \quad (3.3)$$

$$\operatorname{div} u = 0 \quad \text{in } \Omega, \quad (3.4)$$

$$u|_{\partial\Omega} = 0 \quad \text{on } \partial\Omega, \quad (3.5)$$

the reduced Stokes operator is associated to the *reduced Stokes resolvent equations*

$$(\lambda - \Delta + \nabla K_N T)u = f \quad \text{in } \Omega, \quad (3.6)$$

$$u|_{\partial\Omega} = 0 \quad \text{on } \partial\Omega. \quad (3.7)$$

The reduced system (3.6)-(3.7) is obtained from (3.3)-(3.5) by expressing the pressure  $p$  in terms of the data  $f$  and the velocity field  $u$ , which goes back to the work of Grubb and Solonnikov [20, Section 4 and 5], cf. [2, Section 3] for details.

The construction in Section 4 is based on the following lemma.

**Lemma 3.2** *Let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , be an exterior domain with  $C^{1,1}$ -boundary, let  $1 < q < \infty$ , and assume that  $(\lambda + A_{0,q})^{-1}$  exists for some  $\lambda \in \mathbb{C} \setminus (-\infty, 0)$ . Then  $(\lambda + A_q)^{-1}$  exists and*

$$A_{0,q}|_{L_\sigma^q(\Omega)} = A_q, \quad (\lambda + A_{0,q})^{-1}|_{L_\sigma^q(\Omega)} = (\lambda + A_q)^{-1}. \quad (3.8)$$

**Proof:** The first statement can be seen as follows: If  $u \in \mathcal{D}(A_{0,q}) \cap L_\sigma^q(\Omega)$ , then  $\operatorname{div}(-\Delta u + \nabla K_N T u) = 0$  in the sense of distributions and

$$\gamma_\nu(-\Delta u + \nabla K_N T u) = -\gamma_\nu \Delta u + \partial_\nu K_N T u|_{\partial\Omega} = 0$$

in the sense of (2.2). Hence  $-\Delta u = (-\Delta + \nabla K_N T)u - \nabla K_N T u$  is the Helmholtz decomposition of  $-\Delta u$ , i.e.,  $(-\Delta + \nabla K_N T)u = P_q(-\Delta)u = A_q u$ .

In order to prove the second relation let  $u = (\lambda + A_{0,q})^{-1}f$  with  $f \in L_\sigma^q(\Omega)$ . Then applying  $\operatorname{div}$  and  $\gamma_\nu$  to (3.6) we conclude for  $g = \operatorname{div} u$

$$(\lambda - \Delta)g = 0 \quad \text{in } \Omega, \quad (3.9)$$

$$\partial_\nu g = 0 \quad \text{on } \partial\Omega, \quad (3.10)$$

where  $\partial_\nu g|_{\partial\Omega} = \gamma_\nu \nabla g$ . Because of Proposition 3.3 below,  $g = \operatorname{div} u = 0$ . Therefore  $u \in L_\sigma^q(\Omega)$  and  $(\lambda + A_q)u = (\lambda + A_{0,q})u = f$ . Since by the first statement  $\lambda + A_q = (\lambda + A_{q,0})|_{L_\sigma^q(\Omega)}$  is injective, we finally conclude that  $(\lambda + A_q)^{-1}f = u = (\lambda + A_{q,0})^{-1}f$  for every  $f \in L_\sigma^q(\Omega)$ .  $\blacksquare$

It remains to prove:

**Proposition 3.3** *Let  $1 < q < \infty$  and let  $\lambda \in \mathbb{C} \setminus (-\infty, 0)$ . Then there is only one solution  $g \in W_q^1(\Omega)$  with  $g = \operatorname{div} u$  for some  $u \in \mathcal{D}(A_{0,q})$  satisfying (3.9)-(3.10) namely  $g = 0$ .*

**Proof:** If  $\lambda = 0$ , this is a consequence of the (unique) Helmholtz decomposition, cf. [23]. Hence it remains to consider  $\lambda \neq 0$ . Then by definition of  $\gamma_\nu$ , the system (3.9)-(3.10) is equivalent to

$$\lambda(g, v) + (\nabla g, \nabla v) = 0 \quad \text{for all } v \in W_{q'}^1(\Omega). \quad (3.11)$$

If  $q = 2$ , this implies  $g = 0$  by choosing  $v = g$ . Hence in the case  $q \neq 2$  it is sufficient to prove  $g \in W_2^1(\Omega)$ . If  $q > 2$ , let  $\psi \in C^\infty(\mathbb{R}^n)$  such that  $\psi(x) = 1$  for  $|x| \geq R + 1$  and  $\psi(x) = 0$  for  $|x| \leq R$ , where  $R > 0$  is chosen such that  $\partial\Omega \subset B_R(0)$ . Then  $\tilde{g} := \psi g \in W_q^1(\mathbb{R}^n)$  solves

$$(\lambda - \Delta)\tilde{g} = -2(\nabla\psi) \cdot \nabla\tilde{g} - (\Delta\psi)\tilde{g} =: f \quad \text{in } \mathbb{R}^n,$$

where  $f \in L^q(\mathbb{R}^n)$  has compact support. Therefore  $\tilde{g} \in W_r^2(\mathbb{R}^n)$  for every  $1 < r \leq q$ , which implies  $g \in W_2^1(\Omega)$  if  $q > 2$ . Finally, (3.9)-(3.10) imply that  $\Delta g = \lambda g$  and  $\partial_\nu g|_{\partial\Omega} = 0$ . Hence the same procedure of the elliptic regularity theory as in the proof of [15, Lemma 5.4] yields  $g \in W_q^2(\Omega) \leftrightarrow W_{q_1}^1(\Omega)$ , where  $\frac{1}{q_1} = \frac{1}{q} - \frac{1}{n}$ . Thus in the case  $q_1 \geq 2$  we conclude  $g \in W_2^1(\Omega)$ . If  $q_1 < 2$ , we repeat this argument finitely many times until  $g \in W_{q_m}^1(\Omega)$  with  $q_m > 2$ .  $\blacksquare$

## 4 Analysis of the Resolvent for Large $\lambda$

In order to estimate (1.3) with  $\Gamma$  replaced by  $\Gamma_R := \Gamma \setminus \overline{B_R(0)}$ ,  $R > 0$ , it is sufficient to construct an approximate resolvent  $R_\lambda$  satisfying

$$(\lambda + A_{0,q})^{-1} = R_\lambda + S_\lambda,$$

where  $\|S_\lambda\|_{\mathcal{L}(L^q(\Omega))} \leq C_{q,\delta}(1 + |\lambda|)^{-1-\varepsilon}$  for some  $\varepsilon > 0$  and

$$\left\| \int_{\Gamma_R} h(-\lambda) R_\lambda d\lambda \right\|_{\mathcal{L}(L^q(\Omega))} \leq C_{q,\delta} \|h\|_\infty$$

for all  $h \in H(\delta)$ . The operator  $R_\lambda$  can be constructed using the calculus of pseudodifferential boundary value problems developed by Grubb [19] in a version with non-smooth coefficients, cf. [3, 1]. This approach was already used in [3] to prove the existence of a bounded  $H_\infty$ -calculus in asymptotically flat layers. Since this construction is mainly based on localization and a similar approximation of  $(\lambda + A_{0,q})^{-1}$  in a curved half-space  $\mathbb{R}_\gamma^n = \{(x', x_n) \in \mathbb{R}^n : x_n > \gamma(x')\}$ ,  $\gamma \in C^{1,1}(\mathbb{R}^{n-1})$ , it can easily be modified to the case of an exterior domain – as well as many other classes of domains.

For the present contribution it is not necessary to recall the precise construction of the approximate resolvent  $R_\lambda$  in the curved half-space and all the operators belonging to calculus of pseudodifferential boundary value problems with Hölder-continuous coefficients. We refer to [3] for the details. For the following analysis, it is sufficient to recall the following theorem, which summarizes results obtained in [3].

**Theorem 4.1** *Let  $\mathbb{R}_\gamma^n$ ,  $n \geq 2$ ,  $\gamma \in C^{1,1}(\mathbb{R}^{n-1})$  be a curved half-space,  $1 < q < \infty$ , and let  $\delta \in (0, \pi)$ . Then there is a bounded operator  $R_{\gamma,\lambda}: L^q(\mathbb{R}_\gamma^n)^n \rightarrow W_q^2(\mathbb{R}_\gamma^n)^n$ , which is independent of  $q$ , such that*

$$(\lambda - \Delta + \nabla \tilde{K}_{\gamma,N} T) R_{\gamma,\lambda} f = f + S_{\gamma,\lambda} f \quad \text{in } \mathbb{R}_\gamma^n, \quad (4.1)$$

$$R_{\gamma,\lambda} f = 0 \quad \text{on } \partial\mathbb{R}_\gamma^n \quad (4.2)$$

for every  $f \in L^q(\mathbb{R}_\gamma^n)^n$  and  $\lambda \in \mathbb{C} \setminus (-\infty, 0]$ , where  $\|S_{\gamma,\lambda}\|_{\mathcal{L}(L^q(\mathbb{R}_\gamma^n))} \leq C_{q,\delta}(1 + |\lambda|)^{-\varepsilon}$  uniformly in  $\lambda \in \Sigma_\delta$  for some  $\varepsilon > 0$ . Here  $\tilde{K}_{\gamma,N}: W_q^{-\frac{1}{q}}(\partial\mathbb{R}_\gamma^n) \rightarrow W_q^1(\mathbb{R}_\gamma^n)$  is a bounded

operator, which is independent of  $q$ , satisfying

$$\Delta \tilde{K}_{\gamma,N} a = R'_\gamma a \quad \text{in } \mathbb{R}_\gamma^n, \quad (4.3)$$

$$\partial_\nu \tilde{K}_{\gamma,N} a|_{\partial\Omega} = a + S'_\gamma a \quad \text{on } \partial\mathbb{R}_\gamma^n, \quad (4.4)$$

where  $R'_\gamma: W_q^{-\frac{1}{q}-\varepsilon}(\partial\mathbb{R}_\gamma^n) \rightarrow W_{q,0}^{-1}(\mathbb{R}_\gamma^n)$  and  $S'_\gamma: W_q^{-\frac{1}{q}-\varepsilon}(\partial\mathbb{R}_\gamma^n) \rightarrow W_q^{-\frac{1}{q}}(\partial\mathbb{R}_\gamma^n)$  are bounded operators. Moreover, for every  $R > 0$

$$(1 + |\lambda|) \|R_{\gamma,\lambda}\|_{\mathcal{L}(L^q(\mathbb{R}_\gamma^n))} + \|\nabla^2 R_{\gamma,\lambda}\|_{\mathcal{L}(L^q(\mathbb{R}_\gamma^n))} \leq C_{q,\delta}, \quad \lambda \in \Sigma_\delta, \quad (4.5)$$

$$\left\| \int_{\Gamma_R} h(-\lambda) R_{\gamma,\lambda} d\lambda \right\|_{\mathcal{L}(L^q(\mathbb{R}_\gamma^n))} \leq C_{q,\delta} \|h\|_\infty, \quad h \in H(\delta). \quad (4.6)$$

**Proof:** First of all  $\tilde{K}_{\gamma,N} \equiv \tilde{K}_1$  is defined in [3, Section 5.5] as

$$\tilde{K}_{\gamma,N} = F^{*, -1} \underline{k}_1(D_x, x') F_0^*,$$

where  $\underline{k}_1(D_x, x')$  is a Poisson operator of order  $-1$  in R-form with  $C^{0,1}$ -coefficients in the sense of [3, Section 4],  $F: \mathbb{R}_+^n \rightarrow \mathbb{R}_\gamma^n$  is defined by  $F(x) = (x', x_n + \gamma(x'))$ ,  $(F^{*, -1} f)(x) := f(F^{-1}(x))$  is the push-forward of a function  $f: \mathbb{R}_+^n \rightarrow \mathbb{C}$  by  $F$ , and  $(F_0^* a)(y) := a(F_0(y))$  is the pull-back of a function  $a: \partial\mathbb{R}_\gamma^n \rightarrow \mathbb{C}$  by  $F_0 := F|_{\partial\mathbb{R}_+^n}$ . The statements on  $\tilde{K}_{\gamma,N}$  are a consequence of [3, Lemma 5.15] using

$$L^q(\mathbb{R}^{n-1}; L^1(0, b)) \hookrightarrow W_{q,0}^{-1}(\mathbb{R}_+^n), \quad H_q^{-1}(\mathbb{R}^{n-1}; L^q(\mathbb{R}_+)) \hookrightarrow W_{q,0}^{-1}(\mathbb{R}_+^n),$$

and the fact that  $F^*: W_{q,0}^{-1}(\mathbb{R}_+^n) \rightarrow W_{q,0}^{-1}(\mathbb{R}_\gamma^n)$  is an isomorphism.

The operator  $R_{\gamma,\lambda} \equiv R_{0,\lambda}$  is defined in [3, Section 5.6]. Then (4.1)-(4.2) and the estimate of  $S_{\gamma,\lambda}$  is the statement of [3, Lemma 5.17] if  $\nabla \tilde{K}_{\gamma,N} T R_{\gamma,\lambda}$  is replaced by the operator  $\tilde{G}_\lambda$  defined in [3, Lemma 5.17]. However the estimate

$$\left\| \nabla \tilde{K}_{\gamma,N} T R_\lambda - \tilde{G}_\lambda \right\|_{\mathcal{L}(L^q(\mathbb{R}_\gamma^n))} \leq C_{q,\delta} (1 + |\lambda|)^{-\varepsilon}, \quad \lambda \in \Sigma_\delta,$$

for some  $\varepsilon > 0$  is shown in the last part of the proof of [3, Lemma 5.18].

Finally, (4.5) follows from [3, Estimate (5.35)] and (4.6) is a consequence of [3, Theorem 5.13], [3, Lemma 5.14], and [3, Theorem 3.2].  $\blacksquare$

**Remark 4.2** The operator  $\tilde{K}_{\gamma,N}$  of the latter theorem is an approximate Poisson operator to the Laplace equation with Neumann boundary condition on  $\mathbb{R}_\gamma^n$ . As stated above, it is constructed explicitly in [3] and is an operator of the calculus of pseudodifferential boundary value problems necessary to construct the approximate resolvent of the reduced Stokes operator.

We will use the latter theorem to construct an approximate resolvent in an exterior domain  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , with  $C^{1,1}$ -boundary. In order to localize  $\Omega$  let  $U_j \subset \overline{\Omega}$ ,  $j = 0, \dots, m$  be relatively open sets such that

1. each  $U_j \cap \overline{\Omega}$ ,  $j = 1, \dots, m$ , is bounded and coincides (after rotation) with a relatively open subset of  $\overline{\mathbb{R}_{\gamma_j}^n}$ , where  $\mathbb{R}_{\gamma_j}^n$ ,  $\gamma_j \in C^{1,1}(\mathbb{R}^{n-1})$ , is a curved half-space.
2.  $\partial\Omega \subset \bigcup_{j=1}^m U_j$ ,  $U_0 \cap \partial\Omega = \emptyset$ , and  $\overline{\Omega} \subseteq \bigcup_{j=0}^m U_j$ .

Moreover, let  $\varphi_j \in C_{(0)}^\infty(\overline{\Omega})$ ,  $j = 0, \dots, m$ , be a partition of unity on  $\overline{\Omega}$  such that  $\text{supp } \varphi_j \subset U_j$ ,  $j = 0, \dots, m$ . Finally, let  $\psi_j \in C_{(0)}^\infty(\overline{\Omega})$ ,  $j = 0, \dots, m$ , such that  $\psi_j = 1$  on  $\text{supp } \varphi_j$  and again  $\text{supp } \psi_j \subset U_j$ ,  $j = 0, \dots, m$ . Now we define the approximate resolvent  $R_\lambda$  as

$$R_\lambda f = \sum_{j=0}^m \psi_j R_{j,\lambda} \varphi_j f, \quad f \in L^q(\Omega)^n,$$

where  $R_{j,\lambda} = R_{\gamma_j,\lambda}$ ,  $j = 1, \dots, m$ , is the approximate resolvent on  $\mathbb{R}_{\gamma_j}^n$  due to Theorem 4.1 and  $R_{0,\lambda} = (\lambda - \Delta_{\mathbb{R}^n})^{-1}$  is the resolvent of the Laplace operator on  $\mathbb{R}^n$ . Moreover, we define the approximate Poisson operator

$$\tilde{K}_N a = \sum_{j=1}^m \psi_j \tilde{K}_{\gamma_j,N} \varphi_j a, \quad a \in W_q^{-\frac{1}{q}}(\partial\Omega),$$

where  $\tilde{K}_{\gamma_j,N}$  is the operator due to Theorem 4.1 for  $\mathbb{R}_{\gamma_j}^n$ . Now we have

**Lemma 4.3** *Let  $1 < q < \infty$ ,  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , be an exterior domain with  $C^{1,1}$ -boundary, let  $\tilde{K}_N$  be as above, and let  $K_N$  be the Poisson operator of the Neumann problem as defined in Section 3. Then there is some  $\varepsilon > 0$  such that*

$$\|\nabla(K_N - \tilde{K}_N)Tu\|_q \leq C_q \|u\|_{2-\varepsilon,q},$$

for all  $u \in W_q^{2-\varepsilon}(\Omega)^n$ .

**Proof:** Let  $f \in L^{q'}(\Omega)^n$  be arbitrary and let  $f = f_0 + \nabla p$ ,  $f_0 \in L_\sigma^{q'}(\Omega)$ ,  $p \in \dot{W}_q^1(\Omega)$ , be its Helmholtz decomposition. By Lemma 2.1  $p = p_1 + p_2$ , where  $p_1 \in W_{q'}^1(\Omega)$  and  $p_2 \in L_{\text{loc}}^{q'}(\overline{\Omega})$  with  $\nabla p_2 \in W_{q'}^1(\Omega)$  and  $\|(p_1, \nabla p_2)\|_{1,q'} \leq C_{q'} \|\nabla p\|_{q'}$ .

Then by (4.3)-(4.4)

$$\begin{aligned} (\nabla(K_N - \tilde{K}_N)Tu, f)_\Omega &= (\nabla(K_N - \tilde{K}_N)Tu, \nabla p)_\Omega \\ &= (\nabla(K_N - \tilde{K}_N)Tu, \nabla p_2)_\Omega + \langle (I - \partial_\nu \tilde{K}_N)Tu, p_1 \rangle_{\partial\Omega} - (\Delta \tilde{K}_N Tu, p_1)_\Omega, \end{aligned}$$

where

$$\begin{aligned} |\langle (I - \partial_\nu \tilde{K}_N)Tu, p_1 \rangle_{\partial\Omega}| &\leq \sum_{j=1}^m |\langle \psi_j S'_{\gamma_j} \varphi_j Tu, p_1 \rangle_{\partial\Omega}| \\ &\leq C \|Tu\|_{W_q^{-\frac{1}{q}-\varepsilon}(\partial\Omega)} \|p_1\|_{1,q'} \leq C \|u\|_{2-\varepsilon,q} \|f\|_{q'} \end{aligned}$$

by (3.2) for  $\varepsilon > 0$  suitably small. Moreover,

$$\begin{aligned} & |(\Delta \tilde{K}_N T u, p_1)_\Omega| \\ & \leq \sum_{j=1}^m |(\psi_j R'_{\gamma_j} \varphi_j T u, p_1)_\Omega| + \sum_{j=1}^m |(2(\nabla \psi_j) \cdot \nabla \tilde{K}_{\gamma_j, N} \varphi_j T u + (\Delta \psi_j) \tilde{K}_{\gamma_j, N} \varphi_j T u, p_1)_\Omega|, \end{aligned}$$

where

$$|(\psi_j R'_{\gamma_j} \varphi_j T u, p_1)_\Omega| \leq C \|R'_{\gamma_j} \varphi_j T u\|_{W_{q,0}^{-1}(\mathbb{R}_{\gamma_j}^n)} \|p_1\|_{1,q'} \leq C \|u\|_{2-\varepsilon,q} \|f\|_{q'}$$

for some  $\varepsilon > 0$ . Since  $p_1 \in W_{q'}^1(\Omega) \hookrightarrow L^{s'}(\Omega)$  for some  $s' > q'$  and  $\tilde{K}_{\gamma_j, N} : W_s^{-\frac{1}{s}}(\partial \mathbb{R}_{\gamma_j}^n) \rightarrow W_s^1(\mathbb{R}_{\gamma_j}^n)$ ,  $\frac{1}{s} + \frac{1}{s'} = 1$ ,

$$\begin{aligned} & |(2(\nabla \psi_j) \cdot \nabla \tilde{K}_{\gamma_j, N} \varphi_j T u + (\Delta \psi_j) \tilde{K}_{\gamma_j, N} \varphi_j T u, p_1)_\Omega| \\ & \leq C \|\varphi_j T u\|_{W_s^{-\frac{1}{s}}(\partial \mathbb{R}_{\gamma_j}^n)} \|p_1\|_{1,q'} \leq C \|T u\|_{W_q^{-\frac{1}{q}-\varepsilon}(\partial \mathbb{R}_{\gamma_j}^n)} \|f\|_{q'} \leq C \|u\|_{2-\varepsilon,q} \|f\|_{q'}, \end{aligned}$$

where we have used that  $\text{supp } \varphi_j$  is compact and  $-\frac{1}{s} \leq -\frac{1}{q} - \varepsilon$  for suitably small  $\varepsilon > 0$ . The term  $(\nabla \tilde{K}_N T u, \nabla p_2)_\Omega$  is estimated in the same way using  $\nabla p_2 \in W_{q'}^1(\Omega)$ . Finally, by (3.1)

$$\begin{aligned} & |(\nabla K_N T u, \nabla p_2)_\Omega| = |((\Delta - \nabla \text{div})u, \nabla p_2)_\Omega| \\ & \leq |(\nabla u, \nabla^2 p_2)_\Omega| + |(\partial_\nu u, \nabla p_2)_{\partial \Omega}| + |(\text{div } u, \Delta p_2)_\Omega| + |(\text{div } u, \partial_\nu p_2)_{\partial \Omega}| \\ & \leq C \left( \|\nabla u\|_q \|\nabla^2 p_2\|_{q'} + \|\nabla u\|_{L^q(\partial \Omega)} \|\nabla p_2\|_{L^{q'}(\partial \Omega)} \right) \\ & \leq C \|u\|_{2-\varepsilon,q} \|\nabla p_2\|_{1,q'} \leq C \|u\|_{2-\varepsilon,q} \|f\|_{q'} \end{aligned}$$

for some  $\varepsilon > 0$ . ■

Using the latter lemma and Theorem 4.1 we obtain:

**Theorem 4.4** *Let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , be an exterior domain with  $C^{1,1}$ -boundary,  $\delta \in (0, \pi)$ , and let  $1 < q < \infty$ . Then for every  $R > 0$*

$$\left\| \int_{\Gamma_R} h(-\lambda)(\lambda + A_q)^{-1} d\lambda \right\|_{\mathcal{L}(L^q(\Omega))} \leq C_{q,\delta,R} \|h\|_\infty$$

for every  $h \in H(\delta)$ .

**Proof:** First of all, by (2.6), (4.5), and interpolation

$$\|R_{j,\lambda} \varphi_j f\|_{s,q} \leq C_{q,\delta,R} (1 + |\lambda|)^{-1+\frac{\delta}{2}} \|f\|_q, \quad \lambda \in \Sigma_\delta, |\lambda| \geq R, \quad (4.7)$$

for all  $s \in [0, 2]$ ,  $f \in L^q(\Omega)^n$ , and  $j = 0, \dots, m$ . Moreover, by (4.1)

$$\begin{aligned} & (\lambda - \Delta + \nabla K_N T) R_\lambda f \\ &= f + \sum_{j=1}^m \psi_j S_{\gamma_j, \lambda} \varphi_j f - \sum_{j=0}^m (2(\nabla \psi_j) \cdot \nabla R_{j, \lambda} \varphi_j f + (\Delta \psi_j) R_{j, \lambda} \varphi_j f) \\ & \quad + (\nabla K_N T - \nabla \tilde{K}_N T) R_\lambda f. \end{aligned}$$

Hence (4.7), Theorem 4.1, and Lemma 4.3 imply

$$(\lambda - \Delta + \nabla K_N T) R_\lambda = I + S'_\lambda,$$

where  $\|S'_\lambda\|_{\mathcal{L}(L^q(\Omega))} \leq C_{q, \delta} (1 + |\lambda|)^{-1-\varepsilon}$  uniformly in  $\lambda \in \Sigma_\delta$  for some  $\varepsilon > 0$ . Therefore  $(\lambda + A_{0, q})^{-1}$  exists for all  $\lambda \in \Sigma_\delta$  with  $|\lambda| \geq R'$  for some  $R' > 0$  and

$$(\lambda + A_{0, q})^{-1} = R_\lambda + S_\lambda,$$

where  $\|S_\lambda\|_{\mathcal{L}(L^q(\Omega))} \leq C_{q, \delta} (1 + |\lambda|)^{-\varepsilon}$  uniformly in  $\lambda \in \Sigma_\delta$ ,  $|\lambda| \geq R$ . Since  $(\lambda + A_{0, q})^{-1}|_{L^q(\Omega)} = (\lambda + A_q)^{-1}$ , we conclude that

$$(\lambda + A_q)^{-1} = R_\lambda|_{L^q(\Omega)} + S_\lambda|_{L^q(\Omega)},$$

for  $\lambda \in \Sigma_\delta$  with  $|\lambda| \geq R'$ . Moreover,

$$\left\| \int_{\Gamma_R} h(-\lambda) B_\lambda d\lambda \right\|_{\mathcal{L}(L^q(\Omega))} \leq C_{q, \delta} \|h\|_\infty \quad \text{for } B_\lambda = R_\lambda, S_\lambda,$$

for  $R \geq R'$  because of (4.6) and  $\|S_\lambda\| \leq C_{q, \delta} (1 + |\lambda|)^{-1-\varepsilon}$ . Finally, since  $(\lambda + A_q)^{-1}$  is uniformly bounded on each compact subset of  $\overline{\Sigma_\delta} \setminus \{0\}$ ,  $R > 0$  can be chosen arbitrarily.  $\blacksquare$

## 5 Analysis of the Resolvent for Small $\lambda$

First of all, recall that the resolvent of the Stokes operator  $A_{q, \mathbb{R}^2}$  on  $L^q(\mathbb{R}^2)$ ,  $1 < q < \infty$ , can be written as

$$\begin{aligned} (\lambda + A_{q, \mathbb{R}^2})^{-1} &= P(\lambda - \Delta_{\mathbb{R}^2})^{-1}, \quad \text{where} \\ P f &= \mathcal{F}^{-1} \left[ \left( I - \frac{\xi \xi^T}{|\xi|^2} \right) \mathcal{F}[f](\xi) \right] \end{aligned} \tag{5.1}$$

for  $\lambda \in \mathbb{C} \setminus (-\infty, 0]$ . Moreover,  $(\lambda - \Delta_{\mathbb{R}^2})^{-1}$  can be represented as

$$(\lambda - \Delta_{\mathbb{R}^2})^{-1} f = e_\lambda * f, \quad \text{where } e_\lambda(x) = K_0(\sqrt{|\lambda|}|x|).$$

Here  $\sqrt{\lambda}$  denotes the unique square root of  $\lambda$  with  $\operatorname{Re} \sqrt{\lambda} > 0$  and  $K_n(z)$ ,  $n \in \mathbb{N}_0$ , is the modified Bessel function of order  $n$ , cf. [5, page 375]. We note that  $K'_0(z) = -K_1(z)$ . From the definition it follows that

$$K_0(z) = O(\ln z), \quad K_1(z) = O(z^{-1}) \quad \text{as } z \rightarrow 0. \quad (5.2)$$

Furthermore, we will use that

$$|K_n(z)| \leq C_{n,\delta} e^{-\alpha|z|} \quad \text{for } z \in \Sigma_{\delta/2}, |z| \geq 1 \quad (5.3)$$

with  $0 < \delta < \pi$  and for some  $\alpha > 0$  depending on  $\delta$ , cf. [5, page 378].

Moreover, it is well-known that

$$\|(\lambda - \Delta_{\mathbb{R}^2})^{-1} f\|_r \leq C_{s,q,\delta} |\lambda|^{-1+\frac{1}{q}-\frac{1}{r}} \|f\|_q, \quad \lambda \in \Sigma_\delta, \quad (5.4)$$

for  $1 < q < \infty$ ,  $q \leq r \leq \infty$ , and  $\delta \in (0, \pi)$ , cf. e.g. proof of [11, Proposition 4.1]. Because of the identity  $(\lambda + A_{q,\mathbb{R}^2})^{-1} = (\lambda - \Delta_{\mathbb{R}^2})^{-1} P$  and the continuity  $P: L^q(\mathbb{R}^2)^2 \rightarrow L^q(\mathbb{R}^2)^2$ , the same is true for the Stokes resolvent.

Finally,  $A_{q,\mathbb{R}^2}$  possesses a bounded  $H_\infty$ -calculus for all  $\delta \in (0, \pi)$ . In particular,

$$\left\| \frac{1}{2\pi i} \int_\Gamma h(-\lambda) (\lambda + A_{q,\mathbb{R}^2})^{-1} d\lambda \right\|_{\mathcal{L}(L^q(\mathbb{R}^2))} \leq C_\delta \|h\|_\infty, \quad h \in H(\delta), \quad (5.5)$$

cf. [13, Theorem 7.2].

**Remark 5.1** The estimate (5.5) also holds with  $\Gamma$  replaced by  $\Gamma_R = \Gamma \setminus B_R(0)$ ,  $R > 0$ , cf. [3, Lemma 5.14].

Next we recall the representation of the resolvent of the Stokes operator developed in [11], which is

$$(\lambda + A_q)^{-1} f = ((\lambda + A_{q,\mathbb{R}^2})^{-1} f_0)|_\Omega - B_\lambda^\dagger K_\lambda^{-1} f_\lambda, \quad (5.6)$$

where  $f_\lambda = (\lambda + A_{q,\mathbb{R}^2})^{-1} f_0|_{\partial\Omega}$  and  $f_0$  denotes the extension of  $f$  by zero to  $\mathbb{R}^2$ . Moreover,

$$B_\lambda^\dagger \phi = D_\lambda^\dagger \phi - \eta E_\lambda^\dagger M \phi + \frac{4\pi\alpha}{\ln \sqrt{\lambda}} E_\lambda^\dagger \phi, \quad \alpha, \eta \in \mathbb{C} \setminus \{0\}, \quad (5.7)$$

and  $K_\lambda = B_\lambda|_{\partial\Omega}$ . Here  $D_\lambda^\dagger$  is a double layer potential, which satisfies

$$\|D_\lambda^\dagger \phi\|_q \leq \begin{cases} C_{q,\delta} |\lambda|^{\frac{1}{2}-\frac{1}{q}} \|\phi\|_{\infty,\partial\Omega} & \text{if } 1 < q < 2, \\ C_{q,\delta} |\ln \lambda| \|\phi\|_{\infty,\partial\Omega} & \text{if } q = 2, \\ C_{q,\delta} \|\phi\|_{\infty,\partial\Omega} & \text{if } q > 2 \end{cases} \quad (5.8)$$

uniformly in  $\lambda \in \Sigma_\delta$ ,  $\delta \in (0, \pi)$ , with  $|\lambda| \leq \frac{1}{2}$  and  $\phi \in C^0(\partial\Omega)^n$  with  $\int_{\partial\Omega} \nu \cdot \phi \, d\sigma = 0$ , cf. [11, Estimate (4.15)]. Moreover,  $E_\lambda^\dagger$  is the single layer potential defined by

$$E_\lambda^\dagger \phi = \int_{\partial\Omega} E_\lambda^{(r,c)}(x-y) \phi(y) \, d\sigma(y), \quad \phi \in C^0(\partial\Omega)^n,$$

where the matrix  $E_\lambda^{(r,c)}(x) = (E_{jk}^\lambda(x))_{j,k=1,2}$  is defined by

$$E_{jk}^\lambda(x) = \frac{1}{2\pi} \left( \delta_{jk} e_1(\sqrt{\lambda}|x|) + \frac{x_j x_k}{|x|^2} e_2(\sqrt{\lambda}|x|) \right)$$

and

$$e_1(\kappa) = K_0(\kappa) + \kappa^{-1} K_1(\kappa) - \kappa^{-2}, \quad e_2(\kappa) = -K_0(\kappa) - 2\kappa^{-1} K_1(\kappa) + 2\kappa^{-2}.$$

Finally,  $M\phi = \phi - \phi_M$ , where  $\phi_M$  denotes the mean-value of  $\phi$  on  $\partial\Omega$ .

Because of [11, Proposition 3.8]  $K_\lambda$  is invertible for all  $\lambda \in \Sigma_\delta$  with  $|\lambda| \leq R$  for some  $R > 0$  and

$$\|K_\lambda^{-1}\|_{\mathcal{L}(C^0(\partial\Omega))} \leq C_\delta, \quad \lambda \in \Sigma_\delta \cap B_R(0). \quad (5.9)$$

In particular, this implies

$$\|D_\lambda^\dagger K_\lambda^{-1} f_\lambda\|_q \leq C |\lambda|^{-1+\varepsilon} \|f\|_q \quad (5.10)$$

by (5.4) and (5.8) for some  $\varepsilon > 0$ , where  $\int_{\partial\Omega} \nu \cdot f_\lambda \, d\sigma = 0$  since  $\operatorname{div} f_\lambda = 0$  and  $K_\lambda^{-1}$  preserves this property, cf. [11, Lemma 3.7]. Hence the latter term corresponds to an absolutely integrable part in (1.3).

As shown in [11, Section 3]

$$\hat{E}_\lambda^{(r,c)}(\xi) := \mathcal{F}[E_\lambda^{(r,c)}](\xi) = \frac{1}{2\pi(\lambda + |\xi|^2)} \left( I - \frac{\xi \xi^T}{|\xi|^2} \right).$$

Therefore the single layer potential  $E_\lambda^\dagger$  can be represented as  $E_\lambda^\dagger = P E_\lambda'$ , where

$$(E_\lambda' \phi)(x) = \int_{\partial\Omega} K_0(\sqrt{\lambda}|x-y|) \phi(y) \, d\sigma(y)$$

and  $P$  defined as above is bounded on  $L^q(\mathbb{R}^2)$ ,  $1 < q < \infty$ , and is independent of  $\lambda$ . Hence using (5.6) and (5.7) in (1.3) and estimating each term separately we can replace  $E_\lambda^\dagger$  by  $E_\lambda'$ .

Summarizing it suffices to estimate (1.3) with  $(\lambda + A_q)^{-1}$  replaced by  $E_\lambda' M K_\lambda^{-1} f_\lambda$  and  $\frac{1}{\ln \sqrt{\lambda}} E_\lambda' K_\lambda^{-1} f_\lambda$ , which is done in the following two lemmata.

**Lemma 5.2** *Let  $1 < q < \infty$ ,  $\delta \in (0, \pi)$ ,  $\varepsilon > 0$ , and let  $E_\lambda' M$  be defined as above. Then*

$$\|E_\lambda' M \phi\|_q \leq \begin{cases} C_{q,\delta} |\lambda|^{\frac{1}{2}-\frac{1}{q}} \|\phi\|_\infty & \text{if } 1 < q < 2, \\ C_{q,\delta,\varepsilon} |\lambda|^{-\varepsilon} \|\phi\|_\infty & \text{if } q \geq 2 \end{cases}$$

for all  $\phi \in C^0(\partial\Omega)$  uniformly in  $\lambda \in \Sigma_\delta \cap B_1(0)$ . In particular, there is an  $a < 1$  and  $R > 0$  such that for all  $f \in L^q(\mathbb{R}^2)^2$

$$\|E'_\lambda M K_\lambda^{-1} f_\lambda\|_q \leq C_{q,\delta} |\lambda|^{-a} \|f\|_q, \quad \lambda \in \Sigma_\delta, |\lambda| \leq R.$$

**Proof:** First of all, since

$$\|E'_\lambda \phi\|_r \leq C_{r,\delta} |\lambda|^{-\frac{1}{r}} \|\phi\|_\infty, \quad \lambda \in \Sigma_\delta, \quad (5.11)$$

for  $1 < r < \infty$ , cf. [11, Equation (4.14)], and  $L^r(\Omega \cap B_R) \hookrightarrow L^q(\Omega \cap B_R)$  if  $r \geq q$ , it is sufficient to estimate the  $L^q(\Omega_R)$ -norm of  $E'_\lambda M \phi$  with  $\Omega_R := \Omega \setminus B_R(0)$ , where  $R > 1$  is chosen such that  $\partial\Omega \subset B_{R-1}(0)$ . Since  $\int_{\partial\Omega} M \phi(y) d\sigma(y) = 0$ ,

$$E'_\lambda M \phi = \int_{\partial\Omega} \left( K_0(\sqrt{\lambda}|x-y|) - K_0(\sqrt{\lambda}|x|) \right) M \phi(y) d\sigma(y).$$

Moreover,

$$\left| K_0(\sqrt{\lambda}|x-y|) - K_0(\sqrt{\lambda}|x|) \right| \leq C \sup_{s \in [|x-y|, |x|]} |K'_0(\sqrt{\lambda}s)| |\lambda|^{\frac{1}{2}} |y|.$$

Using  $K'_0(z) = -K_1(z)$  and (5.2), we conclude

$$\begin{aligned} & \int_{R \leq |x| \leq |\lambda|^{-\frac{1}{2}}} \left| K_0(\sqrt{\lambda}|x-y|) - K_0(\sqrt{\lambda}|x|) \right|^q dx \\ & \leq C \int_{R \leq |x| \leq |\lambda|^{-\frac{1}{2}}} |x|^{-q} dx \leq \begin{cases} C_\delta |\lambda|^{\frac{q}{2}-1} & \text{if } 1 < q < 2, \\ C_\delta |\ln \lambda| & \text{if } q = 2, \\ C_\delta & \text{if } q > 2, \end{cases} \end{aligned}$$

where we have used  $|x-y| \geq \frac{|x|}{R}$ . Similarly, by (5.3)

$$\begin{aligned} & \int_{|x| > |\lambda|^{-\frac{1}{2}}} \left| K_0(\sqrt{\lambda}|x-y|) - K_0(\sqrt{\lambda}|x|) \right|^q dx \\ & \leq C |\lambda|^{\frac{q}{2}} \int_{|x| > |\lambda|^{-\frac{1}{2}}} e^{-c_q |\lambda|^{\frac{1}{2}} |x|} dx = C' |\lambda|^{\frac{q}{2}-1} \end{aligned}$$

with some  $c_q > 0$ . This implies the statement of the theorem.  $\blacksquare$

**Lemma 5.3** *Let  $1 < q < \infty$ ,  $\delta \in (0, \pi)$ , and let  $E'_\lambda K_\lambda^{-1} f_\lambda$  be defined as above. Then*

$$\left\| \int_{\Gamma'_R} h(-\lambda) \frac{1}{\ln \sqrt{\lambda}} E'_\lambda K_\lambda^{-1} f_\lambda d\lambda \right\|_q \leq C_{q,\delta} \|h\|_\infty \|f\|_q \quad (5.12)$$

for all  $h \in H(\delta)$  and  $f \in L^q(\mathbb{R}^2)^2$ , where  $R > 0$  is as in (5.9).

**Proof:** First of all,

$$\|E'_\lambda K_\lambda^{-1} f_\lambda\|_r \leq C |\lambda|^{-1+\frac{1}{q}-\frac{1}{r}} \|f\|_q$$

for all  $q \leq r < \infty$  due to (5.11), (5.4), and (5.9). Hence we can replace the  $L^q(\Omega)$  on the left-hand side of (5.12) by  $L^q(\Omega_{R'})$  with  $\Omega_{R'} := \Omega \setminus B_{R'}(0)$ , where  $R' > 1$  is chosen such that  $\partial\Omega \subset B_{R'-1}(0)$ . Moreover, because of the Marcinkiewisz interpolation theorem, cf. [8, Theorem 1.3.1], and since  $1 < q < \infty$  is arbitrary, it is sufficient to prove (5.12) with the  $L^q(\Omega_{R'})$ -norm replaced by the weak  $L^q$ -norm on  $\Omega_{R'}$ , i.e.,

$$\|f\|_{L_*^q(\Omega_{R'})} := \sup_{\sigma>0} \{\sigma m(f, \sigma)^{\frac{1}{q}}\}, \quad \text{where } m(f, \sigma) = |\{x \in \Omega_{R'} : |f(x)| > \sigma\}|.$$

Moreover,

$$\left| \int_{\Gamma'_R} h(-\lambda) \frac{1}{\ln \sqrt{\lambda}} (E'_\lambda K_\lambda^{-1} f_\lambda)(x) d\lambda \right| \leq \|h\|_\infty \int_{\Gamma'_R} \left| \frac{1}{\ln \sqrt{\lambda}} (E'_\lambda K_\lambda^{-1} f_\lambda)(x) \right| d|\lambda|.$$

Now by the definition of  $E'_\lambda$  and (5.9)

$$\left| \frac{1}{\ln \sqrt{\lambda}} (E'_\lambda K_\lambda^{-1} f_\lambda)(x) \right| \leq C_q \int_{\partial\Omega} |K_0(\sqrt{\lambda}|x-y)| d\sigma(y) \|\ln \lambda|^{-1} f_\lambda\|_{\infty, \partial\Omega}.$$

Setting  $g_\lambda = \int_{\partial\Omega} |K_0(\sqrt{\lambda}|x-y)| d\sigma(y)$ , we conclude

$$\begin{aligned} & \left| \int_{\Gamma'_R} h(-\lambda) \frac{1}{\ln \sqrt{\lambda}} (E_\lambda K_\lambda^{-1} f_\lambda|_{\partial\Omega})(x) d\lambda \right| \\ & \leq C \|h\|_\infty \left( \int_{\Gamma'_R} g_\lambda(x)^q d|\lambda| \right)^{\frac{1}{q}} \left( \int_{\Gamma'_R} |\ln \lambda|^{-q'} \|f_\lambda\|_{\infty, \partial\Omega}^{q'} d|\lambda| \right)^{\frac{1}{q'}}. \end{aligned}$$

Because of (5.4)

$$\int_{\Gamma'_R} |\ln \lambda|^{-q'} \|f_\lambda\|_{\infty, \partial\Omega}^{q'} d|\lambda| \leq C_q \int_{\Gamma'_R} |\ln \lambda|^{-q'} |\lambda|^{-1} d|\lambda| \|f\|_q^{q'} \leq C'_q \|f\|_q^{q'}.$$

Furthermore,

$$\begin{aligned} \int_{\Gamma'_R} g_\lambda(x)^q d|\lambda| & \leq C \sum_{\pm} \sup_{y \in \partial\Omega} \int_0^R \left| K_0 \left( s^{\frac{1}{2}} e^{\pm i \frac{\delta}{2}} |x-y| \right) \right|^q ds \\ & \leq C \sup_{y \in \partial\Omega} \sum_{\pm} \int_0^\infty \left| K_0 \left( t^{\frac{1}{2}} e^{\pm i \frac{\delta}{2}} \right) \right|^q dt |x-y|^{-2} \\ & \leq C'_{q'} (1+|x|)^{-2} \end{aligned}$$

because  $\text{dist}(x, \partial\Omega) \geq 1$  and by (5.2)-(5.3). Hence  $\left( \int_{\Gamma'_R} g_\lambda(x)^q d|\lambda| \right)^{\frac{1}{q}} \in L_*^q(\Omega_R)$ , which finishes the proof.  $\blacksquare$

**Proof of Theorem 1.1:** Let  $R > 0$  be small enough such that  $K_\lambda^{-1}$  exists for  $\lambda \in \Sigma_\delta$ ,  $|\lambda| \leq R$ . By Theorem 4.4 it is sufficient to prove (1.4) with  $\Gamma$  replaced by  $\Gamma'_R = \Gamma \cap B_R(0)$  and arbitrary  $h \in H(\delta)$ . But this is a consequence of the identity (5.6), Remark 5.1, (5.10), Lemma 5.2, and Lemma 5.3. ■

## References

- [1] H. Abels. *Pseudodifferential Boundary Value Problems with Non-Smooth Coefficients*. Preprint No. 2343, TU Darmstadt, 2004, to appear in Comm. Part. Diff. Eq.
- [2] H. Abels. *Reduced and generalized Stokes resolvent equations in asymptotically flat layers, part I: unique solvability*. Preprint No. 2293, TU Darmstadt, 2003, to appear in J. Math. Fluid. Mech.
- [3] H. Abels. *Reduced and generalized Stokes resolvent equations in asymptotically flat layers, part II:  $H_\infty$ -calculus*. Preprint No. 2294, TU Darmstadt, 2003, to appear in J. Math. Fluid. Mech.
- [4] H. Abels. *Bounded imaginary powers of the Stokes operator in an infinite layer*. J. Evol. Eq. 2, 439-457, 2002.
- [5] M. Abramowitz and I. A. Stegun. *Handbook of Mathematical Functions*. New York: Dover, 1970.
- [6] R. A. Adams. *Sobolev Spaces*. Academic Press, New York, San Francisco, London, 1975.
- [7] H. Amann, M. Hieber, and G. Simonett. *Bounded  $H_\infty$ -calculus for elliptic operators*. Diff. Int. Eq., Vol. 7, No. 3, 613-653, 1994.
- [8] J. Bergh and J. Löfström. *Interpolation Spaces*. Springer, Berlin - Heidelberg - New York, 1976.
- [9] W. Borchers and T. Miyakawa. *On some coercive estimates for the Stokes operator in unbounded domains*. In: Heywood et al. (eds.) The Navier-Stokes Equations, Theory and Numerical Methods 2. Proceedings Oberwolfach 1991 (Lect. Notes Math., vol. 1530, pp. 71-84) Berlin Heidelberg New York, Springer, 1992.
- [10] W. Borchers and H. Sohr. *On the semigroup of the Stokes operator for exterior domains in  $L^q$ -spaces*. Math. Z. 196, 415-425, 1987.
- [11] W. Borchers and W. Varnhorn. *On the boundedness of the Stokes semigroup in two-dimensional exterior domains*. Math. Z. 213, 275-299, 1993.

- [12] S.K. Chua. *Extension theorems on weighted Sobolev space*. Indiana Univ. Math. J., 41, 1027-1076, 1992.
- [13] W. Desch, M. Hieber, and J. Prüss.  *$L^p$ -theory of the Stokes equation in a half-space*. J. Evol. Eq. 1, 115-142, 2001.
- [14] G. Dore and A. Venni. *On the closedness of the sum of two closed operators*. Math. Z. 196, 189-201, 1987.
- [15] R. Farwig and H. Sohr. *Generalized resolvent estimates for the Stokes system in bounded and unbounded domains*. J. Math. Soc. Japan 46, No. 4, 607-643, 1994.
- [16] Y. Giga. *Domains of fractional powers of the Stokes operator in  $L_r$  Spaces*. Arch. Ration. Mech. Anal. 89, 251-265, 1985.
- [17] Y. Giga and H. Sohr. *On the Stokes operator in exterior domains*. J. Fac. Sci. Univ. Tokyo Sect. IA Math. 36, 103-130, 1989.
- [18] Y. Giga and H. Sohr. *Abstract  $L^p$  estimates for the Cauchy problem with applications to the Navier-Stokes equations in exterior domains*. J. Funct. Anal. 102, 72-94, 1991.
- [19] G. Grubb. *Functional Calculus of Pseudodifferential Boundary Problems, 2nd Edition*. Birkhäuser, Basel - Boston - Berlin, 1996.
- [20] G. Grubb and V. A. Solonnikov. *Boundary value problems for the nonstationary Navier-Stokes equations treated by pseudo-differential methods*. Math. Scand. 69, 217-290, 1991.
- [21] A. McIntosh. *Operators which have an  $H_\infty$ -calculus*. in "Miniconference on Operator Theory and Partial Differential Equations", B. Jefferies, A. McIntosh, W. Ricker, editors Proc. Center Math. Anal. A.N.U., 14, 210-231, 1986.
- [22] A. Noll and J. Saal.  *$H^\infty$ -calculus for the Stokes operator on  $L_q$ -spaces*. Math. Z. 244, 651-688, 2003.
- [23] C.G. Simader and H. Sohr. *A new approach to the Helmholtz decomposition and the Neumann problem in  $L_q$ -spaces for bounded and exterior domains*. Series on Advances in Mathematics for Applied Sciences, Vol. 11, Singapore, World Scientific, 1-35, 1992.