

# Runge-Kutta methods for third order weak approximation of SDEs with multidimensional additive noise

**Kristian Debrabant**

Technische Universität Darmstadt, Fachbereich Mathematik, Schloßgartenstr.7, D-64289 Darmstadt, Germany  
debrabant@mathematik.tu-darmstadt.de



TECHNISCHE  
UNIVERSITÄT  
DARMSTADT

Research Group  
Numerical Analysis and  
Scientific Computing

---



---

**Abstract**

---

A new class of third order Runge-Kutta methods for stochastic differential equations with additive noise is introduced. In contrast to Platen's method, which to the knowledge of the author has been up to now the only known third order Runge-Kutta scheme for weak approximation, the new class of methods affords less random variable evaluations and is also applicable to SDEs with multidimensional noise. Order conditions up to order three are calculated and coefficients of a four stage third order method are given. This method has deterministic order four and minimized error constants, and needs in addition less function evaluations than the method of Platen. Applied to a simple example, the new method is compared numerically with Platen's method and some well known second order methods and yields very promising results.

---

---

## 1 Introduction

---

In many applications, e. g., in epidemiology and financial mathematics, taking stochastic effects into account when modelling dynamical systems often leads to stochastic differential equations (SDEs). An important subclass of these are SDEs with additive noise in the form

$$X(t) = x_0 + \int_{t_0}^t g_0(s, X(s)) ds + \sum_{i=1}^m g_i(W_i(t) - W_i(t_0)). \quad (1)$$

Here,  $W(t)$  is an  $m$ -dimensional Wiener process defined on a probability space  $(\Omega, \mathcal{A}, \mathcal{P})$ , the Borel-measurable drift  $g_0 : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is assumed to be sufficiently differentiable and to satisfy a Lipschitz and a linear growth condition, and  $g_i \in \mathbb{R}^d$ ,  $i = 1, \dots, m$ . Then the Existence and Uniqueness Theorem [10] applies. Examples of such systems arising in experimental psychology, turbulent diffusion, radio-astronomy and blood clotting dynamics can be found in [11].

In recent years, the development of numerical methods for the approximation of SDEs has become a field of increasing interest, see e. g. [11, 15] and references therein. Whereas strong approximation methods are designed to obtain good pathwise solutions, see e. g. [3], weak approximation focuses on the expectation of functionals of the solution:

Let  $C_p^l(\mathbb{R}^d, \mathbb{R})$  denote the space of all  $g \in C^l(\mathbb{R}^d, \mathbb{R})$  fulfilling a polynomial growth condition [11]. Further, let  $I^h = \{t_0, t_1, \dots, t_N\}$  with  $t_0 < t_1 < \dots < t_N = T$  be a discretization of the time interval  $I = [t_0, T]$  with step sizes  $h_n = t_{n+1} - t_n$  for  $n = 0, 1, \dots, N-1$ .

**Definition 1** (weak convergence). *A time discrete approximation  $Y^h = (Y^h(t))_{t \in I^h}$  converges weakly with order  $p$  to  $X$  as  $h \rightarrow 0$  at time  $t \in I^h$  if for each  $f \in C_p^{2(p+1)}(\mathbb{R}^d, \mathbb{R})$  there exist a constant  $C_f$  and a finite  $\delta_0 > 0$  such that*

$$|\mathbb{E}(f(Y^h(t))) - \mathbb{E}(f(X(t)))| \leq C_f h^p$$

holds for each  $h \in ]0, \delta_0[$ .

Many approximation schemes for SDEs fall into the class of stochastic Runge-Kutta (SRK) methods. Second order SRK methods for the weak approximation of SDEs were proposed by Kloeden and Platen [11], Komori [13], Mackevicius and Navikas [14], Tocino and Vigo-Aguiar [21], Rößler [18, 19] and Debrabant and Rößler [7, 8, 9]. An explicit third order weak SRK method for autonomous SDEs with additive scalar noise as well as its generalization to general scalar noise have been given in Kloeden and Platen [11]. However, the authors state there that "it remains an open and challenging task to derive simpler derivative free order 3.0 weak schemes, at least for important classes of stochastic differential equations." The present article solves this problem in the case of additive noise and overcomes also the restriction to scalar additive noise.

To do so, we consider the following class of  $s$ -stage SRK methods,

$$Y_{n+1} = Y_n + h_n \sum_{i=1}^s \alpha_i g_0(t + c_i h_n, H_i) + \sqrt{h_n} \sum_{l=1}^m g_l J_l, \quad (2a)$$

$$H_i = Y_n + h_n \sum_{j=1}^s a_{ij} g_0(t + c_j h_n, H_j) + \sqrt{h_n} \sum_{l=1}^m g_l (b_{1,i} J_l + b_{2,i} J_{m+1}), \quad (2b)$$

which defines a  $d$ -dimensional approximation process  $Y^h$  with  $Y^h(t_n) = Y_n$ . Here,  $J_k$ ,  $k = 1, \dots, 2m$ , are independent random variables which do not depend on  $h_n$  and whose moments all exist. Further,  $\alpha = (\alpha_1, \dots, \alpha_s)^\top$ ,  $A = (a_{ij})_{i,j=1,\dots,s}$ ,  $c = (c_1, \dots, c_s)^\top$ ,  $b_1 = (b_{1,1}, \dots, b_{1,s})^\top$  and  $b_2 = (b_{2,1}, \dots, b_{2,s})^\top$  are the coefficients of the SRK method. In the following we choose  $c = A \mathbb{1}$  with  $\mathbb{1} = (1, \dots, 1)^\top \in \mathbb{R}^s$ . Consequently, from now on we can assume for the analysis of this methods that SDE (1) is given in autonomous form, i. e.,  $g_0(t, X) \equiv g_0(X)$ . The analysis relies on the theory of stochastic B-series, which is shortly reviewed in Section 2 and applied in Section 3 to derive order conditions for method (2) up to order three. Then, in Section 4 a concrete explicit third order method is constructed by minimizing the error coefficients. Finally, in Section 5 we give a numerical example.

---

## 2 Stochastic B-series

---

Order conditions for method (2) can be calculated using the colored rooted tree theories derived for the weak approximation of Itô respectively Stratonovich SDEs by SRK methods, compare [16, 17, 12]. Here, we will follow the more general approach developed in [6], which is based on the work in [1, 2, 17] and applicable both for Itô- and Stratonovich SDEs as well as strong and weak approximation. For more details and proofs, see [6].

First, we introduce the set of colored, rooted trees related to the SDE (1), as well as the elementary differentials associated with each of these trees. We adapt these definitions to the special case of additive noise by neglecting all terms which are related to derivatives of  $g_l, l = 1, \dots, m$ .

**Definition 2** (trees). *The set of  $m + 1$ -colored, rooted trees*

$$T^{add} = \{\emptyset\} \cup T_0 \cup \{\bullet_1, \dots, \bullet_m\}$$

related to additive noise is recursively defined as follows:

(a) The graph  $\bullet_0 = [\emptyset]_0$  with only one vertex of color 0 belongs to  $T_0$ .

Let  $\tau = [\tau_1, \tau_2, \dots, \tau_\kappa]_0$  be the tree formed by joining the subtrees  $\tau_1, \tau_2, \dots, \tau_\kappa$  each by a single branch to a common root of color 0.

(b) If  $\tau_1, \tau_2, \dots, \tau_\kappa \in T^{add}$ , then  $\tau = [\tau_1, \tau_2, \dots, \tau_\kappa]_0 \in T_0$ .

Thus,  $T_0$  is the set of trees with a 0-colored root.  $\bullet_0$  will be called deterministic node,  $\bullet_l$  for  $l > 0$  stochastic node of color  $l$ .

**Definition 3** (elementary differentials). *For a tree  $\tau \in T^{add}$  the elementary differential is a mapping  $F(\tau) : \mathbb{R}^d \rightarrow \mathbb{R}^d$  defined recursively by*

(a)  $F(\emptyset)(x_0) = x_0$ ,

(b)  $F(\bullet_0)(x_0) = g_0(x_0)$ ,  $F(\bullet_l)(x_0) = g_l$  for  $l = 1, \dots, m$ ,

(c) If  $\tau = [\tau_1, \tau_2, \dots, \tau_\kappa]_0 \in T_0$ , then

$$F(\tau)(x_0) = g_0^{(\kappa)}(x_0) (F(\tau_1)(x_0), F(\tau_2)(x_0), \dots, F(\tau_\kappa)(x_0)).$$

To simplify the presentation, we neglect in the following the index  $n$  of  $h_n$  and write only  $h$ . Further, we denote by  $\Xi$  the set of families of Borel measurable mappings

$$\Xi := \{ \{ \varphi(h) \}_{h \geq 0} : \varphi(h) : \Omega \rightarrow \mathbb{R} \text{ is } \mathcal{A}\text{-}\mathcal{B}\text{-measurable } \forall h \geq 0 \}.$$

Both the solution of (1) and its approximation by method (2) can formally be written in terms of B-series.

**Definition 4** (B-series). *Given a mapping  $\phi : T^{add} \rightarrow \Xi$  satisfying*

$$\phi(\emptyset) \equiv 1 \text{ and } \phi(\tau)(0) = 0, \quad \forall \tau \in T^{add} \setminus \{\emptyset\}.$$

A (stochastic) B-series is then a formal series of the form

$$B(\phi, x_0; h) = \sum_{\tau \in T^{add}} \alpha(\tau) \cdot \phi(\tau)(h) \cdot F(\tau)(x_0),$$

where  $\alpha : T^{add} \rightarrow \mathbb{Q}$  is given by

$$\alpha(\emptyset) = 1, \quad \alpha(\bullet_l) = 1, \quad \alpha(\tau = [\tau_1, \dots, \tau_\kappa]_l) = \frac{1}{r_1! r_2! \dots r_\kappa!} \prod_{j=1}^{\kappa} \alpha(\tau_j),$$

where  $r_1, r_2, \dots, r_\kappa$  count equal trees among  $\tau_1, \tau_2, \dots, \tau_\kappa$ .

For multidimensional  $\phi : T^{add} \rightarrow \Xi^s, s \in \mathbb{N}$ , we define

$$B(\phi, x_0; h) = [B(\phi_1, x_0; h), \dots, B(\phi_s, x_0; h)]^\top.$$

If  $Z(h)$  can be written as a B-series, then  $f(Z(h))$  can be written as a similar series, where the sum is taken over trees with a root of color  $f$  and subtrees in  $T^{add}$ :

**Lemma 1.** If  $Z(h) = B(\phi, x_0; h)$  is some B-series and  $f \in C^\infty(\mathbb{R}^d, \mathbb{R}^d)$ , then  $f(Z(h))$  can be written as a formal series of the form

$$f(Z(h)) = \sum_{u \in U_f^{add}} \beta(u) \cdot \psi_\phi(u)(h) \cdot G(u)(x_0), \quad (3)$$

where  $U_f^{add}$  is a set of trees derived from  $T^{add}$ , by

- (a)  $[\emptyset]_f \in U_f^{add}$ , and if  $\tau_1, \tau_2, \dots, \tau_\kappa \in T^{add}$ , then  $[\tau_1, \tau_2, \dots, \tau_\kappa]_f \in U_f^{add}$ .
- (b)  $G([\emptyset]_f)(x_0) = f(x_0)$  and  $G(u = [\tau_1, \dots, \tau_\kappa]_f)(x_0) = f^{(\kappa)}(x_0)(F(\tau_1)(x_0), \dots, F(\tau_\kappa)(x_0))$ .
- (c)  $\beta([\emptyset]_f) = 1$  and  $\beta(u = [\tau_1, \dots, \tau_\kappa]_f) = \frac{1}{r_1! r_2! \dots r_q!} \prod_{j=1}^\kappa \alpha(\tau_j)$ , where  $r_1, r_2, \dots, r_q$  count equal trees among  $\tau_1, \tau_2, \dots, \tau_\kappa$ .
- (d)  $\psi_\phi([\emptyset]_f) \equiv 1$  and  $\psi_\phi(u = [\tau_1, \dots, \tau_\kappa]_f)(h) = \prod_{j=1}^\kappa \phi(\tau_j)(h)$ .

**Remark 1.** To simplify the presentation, we assume throughout this article that all derivatives of  $f$  and  $g_0$  exist. Otherwise, one had to consider truncated B-series with a remainder term.

**Theorem 2.** The solution  $X(t_0 + h)$  of (1) can be written as a B-series  $B(\varphi, x_0; h)$  with

$$\begin{aligned} \varphi(\emptyset) &\equiv 1, & \varphi(\bullet_l) &= h, & \varphi(\bullet_l) &= W_l(h), \quad l = 1, \dots, m, \\ \varphi(\tau = [\tau_1, \dots, \tau_\kappa]_0) &= \int_0^h \prod_{j=1}^\kappa \varphi(\tau_j)(s) ds. \end{aligned}$$

The following definition of the order of the tree,  $\rho(\tau)$ , is motivated by the fact that  $E W_l(h)^2 = h$  for  $l \geq 1$ .

**Definition 5 (order).** The order of a tree  $\tau \in T^{add}$  respectively  $u \in U_f^{add}$  is defined by

$$\rho(\emptyset) = 0, \quad \rho(\bullet_l) = \frac{1}{2}, \quad l = 1, \dots, m, \quad \rho(u = [\tau_1, \dots, \tau_\kappa]_f) = \sum_{i=1}^\kappa \rho(\tau_i)$$

and

$$\rho(\tau = [\tau_1, \dots, \tau_\kappa]_l) = \sum_{i=1}^\kappa \rho(\tau_i) + \begin{cases} 1 & \text{for } l = 0, \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

In the following we define the product of vectors by componentwise multiplication.

**Theorem 3.** The numerical approximation  $Y_1$  as well as the stage values can be written in terms of B-series

$$H = B(\Phi_H, x_0; h), \quad Y_1 = B(\Phi, x_0; h)$$

with

$$\Phi_H(\emptyset) \equiv \mathbb{1}, \quad \Phi_H(\bullet_l) = \sqrt{h}(b_1 J_l + b_2 J_{m+l}), \quad l = 1, \dots, m, \quad (4a)$$

$$\Phi_H(\tau = [\tau_1, \dots, \tau_\kappa]_0) = hA \prod_{j=1}^\kappa \Phi_H(\tau_j)(h) \quad (4b)$$

and

$$\Phi(\emptyset) \equiv 1, \quad \Phi(\bullet_l) = \sqrt{h}J_l, \quad l = 1, \dots, m, \quad (5a)$$

$$\Phi(\tau = [\tau_1, \dots, \tau_\kappa]_0) = h\alpha^\top \prod_{j=1}^\kappa \Phi_H(\tau_j)(h). \quad (5b)$$

To determine the weak order we will also need the B-series of the function  $f$ , evaluated at the exact solution and the numerical approximation. From Theorems 2 and 3 and Lemma 1 we obtain

$$f(X(t_0 + h)) = \sum_{u \in U_f^{add}} \beta(u) \cdot \psi_\phi(u)(h) \cdot G(u)(x_0), \quad (6)$$

$$f(Y_1) = \sum_{u \in U_f^{add}} \beta(u) \cdot \psi_{\Phi}(u)(h) \cdot G(u)(x_0), \quad (7)$$

with

$$\psi_{\varphi}([\emptyset]_f) \equiv 1, \quad \psi_{\varphi}(u = [\tau_1, \dots, \tau_{\kappa}]_f)(h) = \prod_{j=1}^{\kappa} \varphi(\tau_j)(h) \quad (8)$$

and

$$\psi_{\Phi}([\emptyset]_f) \equiv 1, \quad \psi_{\Phi}(u = [\tau_1, \dots, \tau_{\kappa}]_f)(h) = \prod_{j=1}^{\kappa} \Phi(\tau_j)(h). \quad (9)$$

---

### 3 Derivation of order conditions

---

With all the B-series in place, we can now present the order conditions for the weak convergence.

Let  $le_f(h; t, x)$  be the weak local error of the method starting at the point  $(t, x)$  with respect to the functional  $f$  and step size  $h$ , i. e.

$$le_f(h; t, x) = \mathbb{E}(f(Y^h(t+h)) - f(X(t+h)) | Y^h(t) = X(t) = x).$$

Inserting (6) and (7), it follows

$$le_f(h; t, x) = \sum_{u \in U_f^{add}} \beta(u) \cdot \mathbb{E}[\psi_{\Phi}(u)(h) - \psi_{\varphi}(u)(h)] \cdot G(u)(x). \quad (10)$$

Thus, we have weak consistency of order  $p$  (and thus, due to the Milstein theorem [15], also weak convergence) if and only if

$$\mathbb{E} \psi_{\Phi}(u)(h) = \mathbb{E} \psi_{\varphi}(u)(h) + \mathcal{O}(h^{p+1}) \quad \forall u \in U_f^{add} \text{ with } \rho(u) \leq p + \frac{1}{2}. \quad (11)$$

Note that (11) slightly weakens conditions given in [17].

By Theorems 2 and 3, (8) and (9) we can now evaluate the order conditions (11) and obtain the following theorem.

**Theorem 4.** *For a  $p$ -th order method choose the independent random variables  $J_k$  of the SRK method (2) such that their moments coincide with those of  $N(0, 1)$  up to the  $(2p+1)$ -th moment for  $k=1, \dots, m$  and up to the  $(2p-1)$ -th moment for  $k=m+1, \dots, 2m$ . If in addition the coefficients of the SRK method (2) fulfill*

1.  $\alpha^\top \mathbf{1} = 1,$

then the method is of weak order  $p = 1$ . If also the equations

2.  $\alpha^\top A \mathbf{1} = \frac{1}{2},$

3.  $\alpha^\top (b_1^2 + b_2^2) = \frac{1}{2},$

4.  $\alpha^\top b_1 = \frac{1}{2}$

are fulfilled, then the SRK method is of weak order  $p = 2$ . Finally, if additionally

5.  $\alpha^\top A^2 \mathbf{1} = \frac{1}{6},$

6.  $\alpha^\top (A \mathbf{1})^2 = \frac{1}{3},$

7.  $\alpha^\top A (b_1^2 + b_2^2) = \frac{1}{6},$

8.  $\alpha^\top (b_1 (Ab_1) + b_2 (Ab_2)) = \frac{1}{6},$

9.  $\alpha^\top Ab_1 = \frac{1}{6},$

10.  $\alpha^\top ((A \mathbf{1}) (b_1^2 + b_2^2)) = \frac{1}{3},$

11.  $\alpha^\top ((A \mathbf{1}) b_1) = \frac{1}{3},$

12.  $\alpha^\top (b_1^2 + b_2^2)^2 = \frac{1}{3},$





13.  $\alpha^\top (b_1^3 + b_1 b_2^2) = \frac{1}{3},$

14.  $\alpha^\top b_1^2 = \frac{1}{3},$

15.  $(\alpha^\top b_2)^2 = \frac{1}{12}$

are fulfilled, then the SRK method is of weak order  $p = 3$ .

Table 1: Relevant second order trees and derivation of corresponding order conditions

$u$	$\Psi_{\Phi}(u)$ $E \Psi_{\Phi}(u)$	$\Psi_{\varphi}(u)$ $E \Psi_{\varphi}(u)$	ord. cond.
	$h^2 J_j^4$ $3h^2$	$W_j(h)^4$ $3h^2$	by assumption
	$h^2 \alpha^\top A \mathbf{1}$ $h^2 \alpha^\top A \mathbf{1}$	$\int_0^h s ds$ $\frac{h^2}{2}$	2.
	$h^2 \alpha^\top (b_1 J_j + b_2 J_{m+j})^2$ $h^2 \alpha^\top (b_1^2 + b_2^2)$	$\int_0^h W_j(s)^2 ds$ $\frac{h^2}{2}$	3.
	$h^2 J_j \alpha^\top (b_1 J_j + b_2 J_{m+j})$ $h^2 \alpha^\top b_1$	$W_j(h) \int_0^h W_j(s) ds$ $\frac{h^2}{2}$	4.

*Proof.* First, we note that  $E \Psi_{\varphi}(u) = 0$  for all trees  $u \in U_f^{add}$  which have an odd number of stochastic nodes of one color, see [5] or also [4, 16]. For those of these trees which have an order  $\rho(u) \leq p + \frac{1}{2}$ , by construction of the method and due to the assumptions on  $J_k$ ,  $k = 1, \dots, 2m$ , it holds also  $E \Psi_{\Phi}(u) = 0$ . Thus, in the following we only have to consider trees with an even number of each kind of stochastic nodes, in particular only trees of integer order. Consequently, there are only two kinds of trees of order one to consider:

$$u_1 = \begin{array}{c} \textcircled{j} \quad \textcircled{j} \\ \diagdown \quad \diagup \\ \textcircled{f} \end{array}, \quad j = 1, \dots, m, \quad \text{and} \quad u_2 = \begin{array}{c} \textcircled{0} \\ | \\ \textcircled{f} \end{array}.$$

Theorems 2 and 3, (8) and (9) yield

$$\Psi_{\Phi}(u_1) = hJ_j^2, \quad \Psi_{\varphi}(u_1) = W_j(h)^2, \quad \Psi_{\Phi}(u_2) = h\alpha^\top \mathbf{1} \quad \text{and} \quad \Psi_{\varphi}(u_2) = h.$$

Thus, by the assumptions on  $J_j$ ,  $E \Psi_{\Phi}(u_1)(h) = E \Psi_{\varphi}(u_1)(h)$  is fulfilled automatically, whereas  $E \Psi_{\Phi}(u_2)(h) = E \Psi_{\varphi}(u_2)(h)$  yields order condition 1.

If  $u \in U_f^{add}$  with  $u = [\tau_1, \dots, \tau_{\kappa}]_f$  can be split into two trees  $u_1 = [\tau_{i_1}, \dots, \tau_{i_{\kappa_1}}]_f$ ,  $u_2 = [\tau_{j_1}, \dots, \tau_{j_{\kappa_2}}]_f$  with disjoint stochastic nodes, i. e. such that  $\kappa_1, \kappa_2 > 0$ ,  $\kappa_1 + \kappa_2 = \kappa$ ,  $\{i_1, \dots, i_{\kappa_1}, j_1, \dots, j_{\kappa_2}\} = \{1, \dots, \kappa\}$  and the sets of colors of the stochastic nodes of  $u_1$  and  $u_2$  are disjoint, then





$$E \Psi_{\Phi}(u)(h) = E \Psi_{\Phi}(u_1)(h) E \Psi_{\Phi}(u_2)(h) = E \Psi_{\varphi}(u_1)(h) E \Psi_{\varphi}(u_2)(h) = E \Psi_{\varphi}(u)(h),$$

provided that the order conditions of orders lower than  $\rho(u)$  are fulfilled. Thus, in the following we only have to consider trees of second and third order which cannot be decomposed into two trees with disjoint stochastic nodes. The relevant second order trees together with the derivation of the corresponding order conditions are given in Table 1, the ones of order three in Tables 2a-2c, which completes the proof.  $\square$

Possible discrete choices for the random variables  $J_k$ ,  $k = 1, \dots, 2m$ , can be found in Table 3.



Table 2a: Relevant third order trees and derivation of corresponding order conditions, part 1

$u$	$\Psi_{\Phi}(u)$ $E \Psi_{\Phi}(u)$	$\Psi_{\varphi}(u)$ $E \Psi_{\varphi}(u)$	ord. cond.
	$h^3 J_j^6$ $15h^2$	$W_j(h)^6$ $15h^2$	by assumption
	$h^3 \alpha^T A^2 \mathbb{1}$ $h^3 \alpha^T A^2 \mathbb{1}$	$\int_0^h \int_0^{s_1} s_2 ds_2 ds_1$ $\frac{h^3}{6}$	5.
	$h^3 \alpha^T (A \mathbb{1})^2$ $h^3 \alpha^T (A \mathbb{1})^2$	$\int_0^h s^2 ds$ $\frac{h^3}{3}$	6.
	$h^3 \alpha^T A (b_1 J_j + b_2 J_{m+j})^2$ $h^3 \alpha^T A (b_1^2 + b_2^2)$	$\int_0^h \int_0^{s_1} W_j(s_2)^2 ds_2 ds_1$ $\frac{h^3}{6}$	7.

#### 4 A concrete explicit third order SRK method

Based on Theorem 4, we now calculate the coefficients of an explicit third order SRK method. The coefficients will be arranged in an extended Butcher array of the form

$$\begin{array}{c|c|c|c} c & A & b_1 & b_2 \\ \hline & \alpha^T & & \end{array}.$$

Whereas in the deterministic case we would only need three stages to construct an explicit third order method, here we need four stages to fulfill the 15 order conditions of Theorem 4. Therefore, we consider  $s = 4$  in (2), but require in addition that the method fulfills also the deterministic order four conditions. The remaining degrees of freedom are then eliminated by minimizing the vector  $lec$  of the order four coefficients of the local error (10) in the Euclidean norm assuming two dimensional noise ( $m = 2$ ), i. e. by minimizing  $\|lec\|_2$  where

$$lec = (\beta(u) \cdot E [\Psi_{\Phi}(u)(h) - \Psi_{\varphi}(u)(h)])_{u \in U_f^{add}, \rho(u)=4}.$$

Using again the B-series analysis, a tedious calculation (one obtains 52 non automatically vanishing terms) and a subsequent attempt of numerical optimization yield the scheme AN3D1 presented in Table 4. AN3D1 needs two random variable and four drift evaluations per step, and thus two random variable and three drift evaluations less than Platen's third order method.

#### 5 Numerical example

In the following, the SRK scheme AN3D1 presented in the last section is applied to compute the second moment of the solution of a simple test equation,

$$X(t) = \frac{1}{10} + \frac{3}{2} \int_0^t X(s) ds + \frac{1}{10} W(t), \quad t \in I = [0, 2]. \quad (12)$$

Table 2b: Relevant third order trees and derivation of corresponding order conditions, part 2






$u$	$\Psi_{\Phi}(u)$ $E \Psi_{\Phi}(u)$	$\Psi_{\varphi}(u)$ $E \Psi_{\varphi}(u)$	ord. cond.
	$h^3 \alpha^T ((b_1 J_j + b_2 J_{m+j})(A(b_1 J_j + b_2 J_{m+j})))$ $h^3 \alpha^T (b_1 (A b_1) + b_2 (A b_2))$	$\int_0^h W_j(s_1) \int_0^{s_1} W_j(s_2) ds_2 ds_1$ $\frac{h^3}{6}$	8.
	$h^3 J_j \alpha^T A (b_1 J_j + b_2 J_{m+j})$ $h^3 \alpha^T A b_1$	$W_j(h) \int_0^h \int_0^{s_1} W_j(s_2) ds_2 ds_1$ $\frac{h^3}{6}$	9.
	$h^3 \alpha^T ((A \mathbb{1})(b_1 J_j + b_2 J_{m+j})^2)$ $h^3 \alpha^T ((A \mathbb{1})(b_1^2 + b_2^2))$	$\int_0^h s W_j(s)^2 ds$ $\frac{h^3}{3}$	10.
	$h^3 J_j \alpha^T ((A \mathbb{1})(b_1 J_j + b_2 J_{m+j}))$ $h^3 \alpha^T ((A \mathbb{1}) b_1)$	$W_j(h) \int_0^h s W_j(s) ds$ $\frac{h^3}{3}$	11.
	$h^3 \alpha^T ((b_1 J_j + b_2 J_{m+j})^2 (b_1 J_k + b_2 J_{m+k})^2)$ $\begin{cases} h^3 \alpha^T (3b_1^4 + 6b_1^2 b_2^2 + 3b_2^4) & j = k \\ h^3 \alpha^T (b_1^2 + b_2^2)^2 & j \neq k \end{cases}$	$\int_0^h W_j(s)^2 W_k(s)^2 ds$ $\begin{cases} h^3 & j = k \\ \frac{h^3}{3} & j \neq k \end{cases}$	12.

Table 2c: Relevant third order trees and derivation of corresponding order conditions, part 3





$u$	$\psi_{\Phi}(u)$ $E \psi_{\Phi}(u)$	$\psi_{\varphi}(u)$ $E \psi_{\varphi}(u)$	ord. cond.
	$h^3 J_k \alpha^{\top} ((b_1 J_j + b_2 J_{m+j})^2 (b_1 J_k + b_2 J_{m+k}))$ $\begin{cases} h^3 \alpha^{\top} (3b_1^3 + 3b_1 b_2^2) & j = k \\ h^3 \alpha^{\top} (b_1^3 + b_1 b_2^2) & j \neq k \end{cases}$	$W_k(h) \int_0^h W_j(s)^2 W_k(s) ds$ $\begin{cases} h^3 & j = k \\ \frac{h^3}{3} & j \neq k \end{cases}$	13.
	$h^3 J_j J_k \alpha^{\top} ((b_1 J_j + b_2 J_{m+j})(b_1 J_k + b_2 J_{m+k}))$ $\begin{cases} h^3 \alpha^{\top} (3b_1^2 + b_2^2) & j = k \\ h^3 \alpha^{\top} b_1^2 & j \neq k \end{cases}$	$W_j(h) W_k(h) \int_0^h W_j(s) W_k(s) ds$ $\begin{cases} \frac{7h^3}{6} & j = k \\ \frac{h^3}{3} & j \neq k \end{cases}$	due to 3. equiv. to 14.
	$h^3 J_j^3 \alpha^{\top} (b_1 J_j + b_2 J_{m+j})$ $3h^3 \alpha^{\top} b_1$	$W_j(h)^3 \int_0^h W_j(s) ds$ $\frac{3h^3}{2}$	4.
	$h^3 (\alpha^{\top} (b_1 J_j + b_2 J_{m+j}))^2$ $h^3 ((\alpha^{\top} b_1)^2 + (\alpha^{\top} b_2)^2)$	$(\int_0^h W_j(s) ds)^2$ $\frac{h^3}{3}$	due to 4. equiv. to 15.

Table 3: Some discrete random variables corresponding up to the  $i$ th moment to  $N(0, 1)$

$i$	distribution
1	$P(J_k = 0) = 1$
3	$P(J_k = 1) = P(J_k = -1) = \frac{1}{2}$
5	$P(J_k = \sqrt{3}) = P(J_k = -\sqrt{3}) = \frac{1}{6}, \quad P(J_k = 0) = \frac{2}{3}$
7	$P(J_k = \sqrt{6}) = P(J_k = -\sqrt{6}) = \frac{1}{30}, \quad P(J_k = 1) = P(J_k = -1) = \frac{3}{10}, \quad P(J_k = 0) = \frac{1}{3}$

Table 4: Coefficients of AN3D1

0	0	0	0	0	$b_1$   $b_2$ with
1	1	0	0	0	
1/2	3/8	1/8	0	0	
1	-0.4526683126055039	-0.4842227708685013	1.9368910834740051	0	
	1/6	-0.005430430675258792	2/3	0.1720970973419255	

$$b_1 = (-0.01844540496323970, 0.8017012756521233, 0.5092227024816198, 0.9758794209767762)^{\top}$$

$$b_2 = (-0.1866426386543421, -0.8575745885712401, -0.4723392695015512, 0.3060354860326548)^{\top}$$




Figure 1: Computational effort per simulation path versus precision

The second moment of the solution can be calculated as

$$E(X^2(t)) = \frac{2}{9} \left( \frac{397}{200} - \frac{23}{5} e^{3/2t} + \frac{133}{50} e^{3t} \right). \quad (13)$$

For this example, we compare the performance of AN3D1 (with  $N(0,1)$ -distributed random variables) with some well known schemes, namely the third and the second order SRK schemes due to Platen [11], denoted here by PL3 and PL2, respectively, DRI1 due to Debrabant and Rößler [9], and the extrapolated Euler-Maruyama scheme EXEM (cp. [20]) also attaining order two, which is given by  $2E\left(\left(Z^{h/2}(t)\right)^2\right) - E\left(\left(Z^h(t)\right)^2\right)$ , based on the Euler-Maruyama approximations  $Z^{h/2}(t)$  and  $Z^h(t)$  calculated with step sizes  $h/2$  and  $h$ .

In each case, the functional  $u(t) = E(X^2(t))$  is approximated by a Monte Carlo simulation. The sample average  $u_{M,h}(t) = \frac{1}{M} \sum_{k=1}^M \left(Y^h(t, \omega_k)\right)^2$ ,  $\omega_k \in \Omega$ , of  $M$  independent simulated realizations of the considered approximation  $Y^h(t)$  is calculated in order to estimate the expectation. In the following, we denote by  $\hat{\mu} = u_{M,h}(T) - u(T)$  the mean error at time  $T$  and by  $\hat{\sigma}_{\mu}^2$  the empirical variance of the mean error. Further, we calculate the confidence interval with boundaries  $a$  and  $b$  to the level of 90% for the estimated error  $\hat{\mu}$  (see [11] for details).

The solution value  $E(X^2(T))$  is approximated with step sizes  $2^1, \dots, 2^{-4}$ , and  $M = 10^9$  simulations are performed in order to determine the systematic error of the considered schemes at time  $T = 2$ . The results for the applied schemes are presented in Table 5. Of course, these results have to be related to the computational effort of the schemes which we take in the following as sum of the number of evaluations of the drift function  $a$  as well as the number of random variables that have to be simulated. Then we can oppose the computational efforts to the errors of the analyzed schemes. The results are presented in Figure 1. Although being of different order, the two Platen schemes yield comparable results. This is due to the much higher computational costs of PL3. Both methods are better than the extrapolated Euler method, but perform worse than DRI1, which has optimized coefficients [9] and behaves therefore nearly like an order three method. Our new method AN3D1 performs best.

---

## 6 Conclusion

---

We have presented a general class of SRK methods for the weak approximation of SDEs with additive noise, together with the corresponding order conditions up to order three. A concrete explicit third order method has been derived, for which a numerical comparison with some well known other methods regarding its performance yielded very promising results. In contrast to the method of Platen, it needs only two random variables and four drift evaluations per step and is also applicable to SDEs driven by a multidimensional Wiener process. Future research may be done by constructing implicit methods with good stability properties, i. e. which are suitable for stiff problems, and by developing methods for more general noise.

---

## Bibliography

---

- [1] Burrage, K., Burrage, P.M.: High strong order explicit Runge–Kutta methods for stochastic ordinary differential equations. Appl. Numer. Math. **22**(1-3), 81–101 (1996). Special issue celebrating the centenary of Runge-Kutta methods

Table 5: Mean errors, empirical variances and confidence intervals

	$h$	$\hat{\mu}$	$\hat{\sigma}_{\mu}^2$	$a$	$b$
EXEM	$2^1$	-1.900E+02	7.882E-07	-1.900E+02	-1.900E+02
	$2^0$	-1.499E+02	7.032E-06	-1.499E+02	-1.499E+02
	$2^{-1}$	-9.357E+01	3.666E-05	-9.357E+01	-9.357E+01
	$2^{-2}$	-4.435E+01	8.881E-05	-4.435E+01	-4.434E+01
	$2^{-3}$	-1.649E+01	1.988E-04	-1.650E+01	-1.649E+01
	$2^{-4}$	-5.170E+00	2.770E-04	-5.174E+00	-5.166E+00
	PL2	$2^1$	-1.840E+02	6.783E-07	-1.840E+02
$2^0$		-1.294E+02	6.348E-06	-1.294E+02	-1.294E+02
$2^{-1}$		-6.412E+01	2.790E-05	-6.412E+01	-6.412E+01
$2^{-2}$		-2.312E+01	4.995E-05	-2.312E+01	-2.312E+01
$2^{-3}$		-6.880E+00	5.861E-05	-6.882E+00	-6.878E+00
$2^{-4}$		-1.863E+00	8.264E-05	-1.865E+00	-1.861E+00
PL3		$2^1$	-8.377E+01	2.936E-03	-8.378E+01
	$2^0$	-2.705E+01	1.614E-03	-2.706E+01	-2.704E+01
	$2^{-1}$	-8.941E+00	2.345E-04	-8.944E+00	-8.937E+00
	$2^{-2}$	-1.951E+00	6.624E-05	-1.953E+00	-1.949E+00
	$2^{-3}$	-3.111E-01	6.180E-05	-3.130E-01	-3.093E-01
	$2^{-4}$	-4.307E-02	4.718E-05	-4.470E-02	-4.144E-02
	DR11	$2^1$	-1.316E+02	3.486E-06	-1.316E+02
$2^0$		-5.438E+01	2.049E-05	-5.438E+01	-5.437E+01
$2^{-1}$		-1.308E+01	4.872E-05	-1.308E+01	-1.308E+01
$2^{-2}$		-2.254E+00	6.097E-05	-2.256E+00	-2.252E+00
$2^{-3}$		-3.314E-01	6.225E-05	-3.333E-01	-3.295E-01
$2^{-4}$		-4.343E-02	8.405E-05	-4.560E-02	-4.126E-02
AN3D1		$2^1$	-7.638E+01	1.286E-05	-7.638E+01
	$2^0$	-1.654E+01	4.729E-05	-1.654E+01	-1.654E+01
	$2^{-1}$	-1.946E+00	6.804E-05	-1.948E+00	-1.944E+00
	$2^{-2}$	-1.651E-01	4.993E-05	-1.668E-01	-1.635E-01
	$2^{-3}$	-1.073E-02	5.940E-05	-1.255E-02	-8.900E-03
	$2^{-4}$	-1.030E-04	4.754E-05	-1.738E-03	1.532E-03

- [2] Burrage, K., Burrage, P.M.: Order conditions of stochastic Runge–Kutta methods by  $B$ -series. *SIAM J. Numer. Anal.* **38**(5), 1626–1646 (electronic) (2000)
- [3] Burrage, K., Burrage, P.M., Tian, T.H.: Numerical methods for strong solutions of stochastic differential equations: an overview. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **460**(2041), 373–402 (2004). Stochastic analysis with applications to mathematical finance
- [4] Burrage, P.M.: Runge–Kutta methods for stochastic differential equations. Ph.D. thesis, The University of Queensland, Brisbane (1999)
- [5] Debrabant, K., Kværnø, A.: Stochastic Taylor expansions: Weight functions of  $B$ -series expressed as multiple integrals. Accepted for publication in *Stoch. Anal. Appl.*
- [6] Debrabant, K., Kværnø, A.:  $B$ -series analysis of stochastic Runge–Kutta methods that use an iterative scheme to compute their internal stage values. *SIAM J. Numer. Anal.* **47**(1), 181–203 (2008/09)
- [7] Debrabant, K., Rößler, A.: Continuous weak approximation for stochastic differential equations. *J. Comput. Appl. Math.* **214**(1), 259–273 (2008)
- [8] Debrabant, K., Rößler, A.: Diagonally drift-implicit Runge–Kutta methods of weak order one and two for Itô SDEs and stability analysis. *Appl. Numer. Math.* **59**(3–4), 595–607 (2009)
- [9] Debrabant, K., Rößler, A.: Families of efficient second order Runge–Kutta methods for the weak approximation of Itô stochastic differential equations. *Appl. Numer. Math.* **59**(3–4), 582–594 (2009)
- [10] Karatzas, I., Shreve, S.E.: Brownian motion and stochastic calculus, *Graduate Texts in Mathematics*, vol. 113, second edn. Springer-Verlag, New York (1991)
- [11] Kloeden, P.E., Platen, E.: Numerical solution of stochastic differential equations, *Applications of Mathematics*, vol. 21, 2 edn. Springer-Verlag, Berlin (1999)
- [12] Komori, Y.: Multi-colored rooted tree analysis of the weak order conditions of a stochastic Runge–Kutta family. *Appl. Numer. Math.* **57**(2), 147–165 (2007)
- [13] Komori, Y.: Weak second-order stochastic Runge–Kutta methods for non-commutative stochastic differential equations. *J. Comput. Appl. Math.* **206**(1), 158–173 (2007)
- [14] Mackevičius, V., Navikas, J.: Second order weak Runge–Kutta type methods of Itô equations. *Math. Comput. Simulation* **57**(1-2), 29–34 (2001)
- [15] Milstein, G.N.: Numerical integration of stochastic differential equations, *Mathematics and its Applications*, vol. 313. Kluwer Academic Publishers Group, Dordrecht (1995). Translated and revised from the 1988 Russian original
- [16] Rößler, A.: Stochastic Taylor expansions for the expectation of functionals of diffusion processes. *Stoch. Anal. Appl.* **22**(6), 1553–1576 (2004)
- [17] Rößler, A.: Rooted tree analysis for order conditions of stochastic Runge–Kutta methods for the weak approximation of stochastic differential equations. *Stoch. Anal. Appl.* **24**(1), 97–134 (2006)
- [18] Rößler, A.: Second order Runge–Kutta methods for Stratonovich stochastic differential equations. *BIT* **47**(3), 657–680 (2007)
- [19] Rößler, A.: Second order Runge–Kutta methods for Itô stochastic differential equations. *SIAM J. Numer. Anal.* **47**(3), 1713–1738 (electronic) (2009)
- [20] Talay, D., Tubaro, L.: Expansion of the global error for numerical schemes solving stochastic differential equations. *Stoch. Anal. Appl.* **8**(4), 94–120 (1990)
- [21] Tocino, A., Vigo-Aguiar, J.: Weak second order conditions for stochastic Runge–Kutta methods. *SIAM J. Sci. Comput.* **24**(2), 507–523 (electronic) (2002)